Agriculture Canada

Publication 1863/E

Beneficial insects and common pests on strawberry and raspberry crops



Canada

91-07128/

Beneficial insects and common pests on strawberry and raspberry crops

Written under contract by
D.E. Henderson
E.S. Cropconsult Ltd.
Vancouver, British Columbia

in collaboration with
D.A. Raworth
Research Station
Vancouver, British Columbia

Cover illustration
Green lacewing searching for aphids (photo by D. Gillespie)

Agriculture Canada Publication 1863/E
available from
Communications Branch, Agriculture Canada
Ottawa, Ont. K1A 0C7

©Minister of Supply and Services Canada 1991
Cat. No. A53-1863/1991E ISBN 0-662-18483-1
Printed 1991 5M-04:91

Produced by Research Program Service

Également disponible en français sous le titre
Insectes utiles et nuisibles du fraisier et du framboisier

Contents

Acknowledgments 6

Introduction 7

Beneficial flies 9
Syrphid flies (Diptera: Syrphidae) 9
Tachinid flies (Diptera: Tachinidae) 9
Aphid midges (Diptera: Cecidomyiidae) 11

Beneficial wasps 11
Ichneumonid wasps (Hymenoptera: Ichneumonidae) 11
Braconid wasps (Hymenoptera: Braconidae) 13
Chalcid wasps (Hymenoptera: Chalcidoidea) 13

Beneficial beetles 14
Coccinellid beetles (Coleoptera: Coccinellidae) 14
Rove beetles (Coleoptera: Staphilinidae) 15
Ground beetles (Coleoptera: Carabidae) 17

Beneficial true bugs 17
Damsel bugs (Hemiptera: Nabidae) 17
Assassin bugs (Hemiptera: Reduviidae) 17
Bigeyed bugs (Hemiptera: Lygaeidae) 20
Minute pirate bugs (Hemiptera: Anthocoridae) 20

Beneficial lacewings 22
Green lacewings (Neuroptera: Chrysopidae) 22
Brown lacewings (Neuroptera: Hemerobiidae) 22

Beneficial mites 23
Predator mites (Acari: Phytoseiidae) 23

Other predators 23

Common insect pests of raspberry and strawberry crops 25
Lygus bug and other plant bugs (Hemiptera: Miridae) 25
Aphids (Homoptera: Aphididae) 25
Leafhoppers and spittlebugs (Homoptera) 26
Weevils and other beetles (Coleoptera) 26
Leafrollers, caterpillars, climbing cutworms, and borers (Lepidoptera) 27
Sawflies (Hymenoptera: Tenthredinidae) 30
Mites (Acari) 30

More information 31

Acknowledgments

This publication was born out of a request by members of the Lower Mainland Horticultural Improvement Association at the annual growers' short course (Clearbrooke, B.C.) in February, 1988. Its production has indeed been a cooperative project. Agriculture Canada's Research Program Service allocated funds and facilities to publish this handbook. The Province of British Columbia, Ministry of Agriculture and Fisheries, Crop Protection Branch provided contract funds and expertise to support the gathering of information and the writing. The following scientists and professionals from across Canada supplied photographs, specimens, and information specific to their region: Dr. R. Smith, Vedalia Research, Galiano, B.C.; Dr. W.G. Wellington and S.G. Cannings, University of British Columbia, Vancouver, B.C.; Dr. D. Gillespie, Agriculture Canada, Agassiz, B.C.; Dr. C. Vincent and Dr. N. Bostanian, Agriculture Canada, Saint-Jean-sur-Richelieu, Que.; Dr. M. Luffman, Agriculture Canada, Bouctouche, N.B.; R. Tremblay, New Brunswick Department of Agriculture, Fredericton, N.B.; Dr. B.D. Schaber, Agriculture Canada, Lethbridge, Alta.; Dr. L. Gilkeson, Applied Bionomics, Sidney, B.C.; Dr. M. Steiner and M. Herbut, Environmental Centre, Vegreville, Alta.; R. Long and Dr. M. Mackauer, Simon Fraser University, Burnaby, B.C.; C. Chan, W. MacDiarmid, H. Severson, and Dr. W.T. Cram, Agriculture Canada, Vancouver, B.C. Editorial comments were provided by the following: Dr. H. Gerber and W. Peters, B.C. Ministry of Agriculture and Fisheries, Abbotsford, B.C., and S. Krannitz, Phero Tech Inc., Delta, B.C. Many thanks to you all.

Introduction

Because of concerns about the environment, many growers are endeavoring to reduce the use of chemicals on their crops, by using other methods to control pests and disease before resorting to chemicals. This approach, called integrated pest management (IPM), is being used more widely each year, both in Canada and elsewhere. To use IPM you must be aware of what other control methods are available and how to use them. This handbook is intended to supply at least some of the information toward this end. Beneficial insects that provide natural control of pest species are very susceptible to pesticides and so, in a field that is sprayed regularly, few survive. For this reason, they are not well known to many growers or even to extension workers. When growers begin using IPM, either individually or with a consultant, they monitor pest populations regularly to determine the threat of damage. Insecticides are used only when an insect pest threatens to increase to the point of causing economic damage. As a result, many insect species increase in number in the field, many of them beneficial. It is important to be able to recognize the beneficial species as well as the pest species. Although several handbooks and publications available to growers across Canada describe pests, none deal with beneficial insects.

This handbook describes beneficial insects found on strawberry and raspberry crops, but many of the same insects are found on other crops with similar pests. For instance, the predators that attack aphids on strawberry will commonly attack aphids on any plant. In addition, whereas some beneficial insects (such as parasitic wasps) may be specific to a single pest species, those that attack other pests commonly appear similar to one described in this book. The handbook, then, will be useful not only for identifying beneficial insects on strawberry and raspberry crops, but also for identifying beneficial insects on almost any crop.

Note to professionals In collecting the information and photographs for this handbook, it became apparent that we lacked information about the presence of beneficial insects and that few photographs were available. Some biological information in this publication is therefore incomplete and a few photographs are missing. If you have information about beneficial insects in your area, and especially if you have good photographs of any of them, we would like to hear from you. We plan to produce similar handbooks for other crops in the future, and contributions of information and photographs will add to their completeness. You can contact either of the authors at the following addresses:

Dr. D.E. Henderson, E.S. Cropconsult Ltd., 2962 West 15th Ave., Vancouver, B.C. V6K 3A3

Dr. D.A. Raworth, Research Station, Research Branch, Agriculture Canada, 6660 N.W. Marine Drive, Vancouver, B.C. V6T 1X2.



Plate 1 ***a*** **Adult syrphid fly. (Photo by W.G. Wellington)**
b **Adult syrphid fly. (Photo by W.G. Wellington)**
c **Syrphid larva. (Photo by C. Chan)**
d **Syrphid larva. (Photo by H. Severson)**
e **Aphid midge adult. (Photo by W. MacDiarmid)**
f **Aphid midge larva attacking an aphid. (Photo by L. Gilkeson)**

Beneficial flies

Syrphid flies (Diptera: Syrphidae)

Syrphids are flies (6–13 mm in length) that have abdominal stripes like wasps and bees but possess no stinger (Plate 1*a* and *b*). The stripes are a warning coloration that protects them against would-be predators. The many species of syrphid flies are pollinators, second only in importance to bees. They feed on pollen, nectar, and "honeydew"—the sticky, shiny substance that contaminates plants around aphid colonies. They are most easily distinguished from bees by their ability to hover like small helicopters in the air. Syrphid or hover fly larvae (the immature stage) feed exclusively on aphids.

Syrphids lay most of their eggs on heavily infested plants among large aphid colonies. Each female produces between 400 and 1000 oblong, white eggs laid singly or in small batches, depending on the species. Within a few days the egg hatches and a minute, slug-like larva emerges, which, as it grows, develops a color pattern characteristic of its species. Some are green with white stripes or markings (Plate 1c), whereas others are yellowish or brownish with a mottled pattern (Plate 1*d*). Larvae search for aphids by raising the front of their body and swinging it from side to side and up and down, searching the leaf surface. When a larva contacts an aphid, it seizes it with its sharp mouthparts, holds it up in the air, and sucks out the body contents. In this way, a single larva can devour 200–800 aphids in its 7–10-day life. When fully grown (up to 6 mm long), the larva seeks out a dark place, such as a dead rolled leaf on the plant, or burrows into the soil; it then pupates (an inactive stage) for about 1 week. The new adult emerges ready to mate and lay more eggs. Syrphids produce several generations a year and are considered to be very effective predators of aphids.

Tachinid flies (Diptera: Tachinidae)

The many species of tachinid flies appear rather like bristly house flies, bees, or wasps and are about the same size (Fig. 1). They are among the most common of parasitic insects. Adults feed on plant secretions as well as honeydew produced by aphids, scales, or leaf hoppers. Larvae, on the other hand, are parasitic within other insects, mainly the immature stages of moths, sawflies, and beetles, but also some plant bugs and grasshoppers. Some specialize in attacking concealed caterpillars such as leaf rollers, whereas others prefer to attack cutworms. Various species have been introduced to North America from Europe to control pests here. One such example is the "cutworm tachina," which is now a common enemy of cutworms in many areas of Canada. Female tachinid flies deposit their eggs directly on hosts, or on foliage that is eaten by their host; some even inject their eggs into the body of a host. Other species produce live young (larvae), which they deposit in similar places as those that lay eggs. White eggs on the body of a caterpillar are a signal that it has been parasitized by a tachinid fly. The newly hatched larva burrows into its host where it feeds



Fig. 1 Tachinid fly parasitic on raspberry looper. (Photo by W.T. Cram)



Fig. 2 Common body shapes of ichneumonid, braconid, and chalcid wasps.

for a number of days on its host's internal organs before leaving to pupate nearby. An insect attacked by a tachinid fly in this way eventually dies.

Aphid midges (Diptera: Cecidomyiidae)

The adult aphid midge (*Aphidoletes aphidimyza* (Rondani)) is 2–3 mm in length and rarely seen because of its nocturnal habits and small, delicate size (Plate 1*e*). Each female lives about 10 days and produces about 250 eggs. With only a meal of plant sap or nectar, a female is able to produce her first batch of about 40 eggs. To produce more than this first batch, however, she must feed on aphid honeydew. Unlike the delicate adults, the tiny (up to 1.5 mm long) orange larvae are tough (Plate 1*f*). They feed on aphids and overwinter out of doors in most parts of Canada. Small larvae capture aphids by puncturing a leg joint and injecting a poison that paralyzes the aphid; the larva then sucks it dry. When aphids are numerous, larvae kill more aphids than they eat. After 1–2 weeks of feeding, the larvae drop to the soil where they burrow and pupate. After 10–14 days, new adults emerge. Experience in British Columbia indicates that aphid midges are most common when aphid numbers are high and are rare when aphid numbers are low. One species is available commercially in Canada and is currently used to control aphids on greenhouse and fruit-tree crops. Some members of this family are also known to attack mite eggs and larvae. Others attack soft-bodied scales and their eggs.

Beneficial wasps

Ichneumonid wasps (Hymenoptera: Ichneumonidae)

Ichneumonid wasps are slender bodied and range in length from 5–36 mm. Females usually have a long, needle-like egg-laying structure (ovipositor) at the end of their abdomen (Fig. 2). The ovipositor is longer than the body of the wasp and is commonly taken to be a "stinger," but its real purpose is to inject eggs into insects, which are the food of the larval stage. Larval forms of ichneumonid wasps are parasitic within the immature stages of other insects, primarily caterpillars of moths and butterflies. The length of the ovipositor allows the female to inject her eggs into hidden hosts such as leaf-rolling or stem-boring caterpillars. An example of such ichneumonids are *Apophua* species, some of which attack strawberry leafrollers (*Ancylis comptana* (Froelich)) while they are hidden inside their leaves (Plate 2*a*). Many species of ichneumonid wasps are common across Canada. Adults are brown, red, or black with variable markings. They can be distinguished from other parasitic wasps by their large size and the length of their abdomen, which is longer than the head and thorax together. The long ovipositor is their other distinguishing feature.



Plate 2 ***a*** **Ichneumonid wasp (*Apophua*) on leafroller cocoon. (Photo by D. Gillespie)**
b **Braconid injecting an egg. (Photo by R. Long)**
c **"Mummies" in an aphid colony. (Photo by M. Mackauer)**
d **Pupating braconid in a mummy. (Photo by M. Mackauer)**
e **Braconid emerging from a mummy (Photo by C. Chan)**
f **Braconid wasp (*Macrocentrus*) cocoons. (Photo by D. Gillespie)**

Braconid wasps (Hymenoptera: Braconidae)

Braconid wasps are smaller than ichneumonids, up to 12 mm in length, and their body shape is shorter and stouter (Fig. 2). They too have an ovipositor but it is carried inside the body until it is needed to inject an egg into a host. Braconids parasitize a broader range of hosts than ichneumonids—from caterpillars, such as the orange tortrix (*Argyrotaenia citrana* (Fernald)), to flies, wasps, beetles, and aphids. After a female injects an egg into a host, the larva feeds slowly on that single host. By the time the host dies, the larva is fully grown. It pupates inside or near the dead host, sometimes in a silken cocoon, to emerge later as an adult wasp. Like the ichneumonid wasps, many species of braconid wasp are found in all parts of Canada.

Braconids that parasitize aphids are an important group of beneficial insects on strawberry and raspberry crops. Many aphid parasites are braconid and their life cycles are similar. The female injects her egg, with lightning speed (Plate 2*b*), into a young aphid while it feeds on the host plant. Usually the aphid is disturbed by the attack but does not leave the plant. A parasitized aphid usually does not produce offspring but continues to feed until the parasite inside completely consumes it. The fully grown larva cements the dead aphid to the leaf surface and the aphid shell becomes parchment-like or black. An aphid killed in such a way is termed a "mummy" and is easy to recognize in a colony of live aphids (Plate 2*c*). The mummy serves as a protective case for the pupating braconid (Plate 2*d*). When fully developed about 1 week later, the adult wasp cuts a round hole in the mummy and emerges (Plate 2*e*). The empty mummy, with its hole, remains on the leaf. The presence of mummies in a colony of aphids is a signal that many of the other healthy-looking aphids may actually be parasitized and will die within days.

Braconids that parasitize caterpillars are another important group. In many cases the life cycle is similar to that of the aphid parasites. A different pattern, however, is found among the *Copidosoma* species. A single egg is laid in the egg of the host. As the host goes through its various stages of development, the wasp egg divides and produces several individual larvae. When the host dies, the larvae pupate within it, to emerge later as adult wasps. In this way the female can produce many offspring even if she only finds one host. Others, *Macrocentrus* species for example, take advantage of a large host by laying many eggs in one caterpillar. The larvae grow and feed, and, when the host dies, wasp pupae can be found clustered in what is left of the host's cocoon (Plate 2*f*).

Chalcid wasps (Hymenoptera: Chalcidoidea)

Most beneficial chalcid wasps are even smaller than the ichneumonids and braconids (Fig. 2). They are up to 7 mm long, their color is commonly dark metallic blue or green, and their wings are clear. As they take flight, they appear to jump. Chalcids are found everywhere but are often overlooked because of their small size. Members of this family of wasps

parasitize caterpillars of moths and butterflies, larvae of some flies, and larvae of some beetles. Some chalcid wasps are not beneficial because they attack tachinid flies and pupae of ichneumonid wasps.

A well-known group of chalcid wasps, *Trichogramma*, lays its eggs in the eggs of many species of moths and butterflies. The adult is less than 1 mm long but is effective as a biological control agent. It is available commercially and is used for a number of crops to control caterpillar pests, less often in Canada than in the United States.

Beneficial beetles

Coccinellid beetles (Coleoptera: Coccinellidae)

Ladybird beetles

Adult ladybird beetles are probably the most easily recognized predators in any agricultural system. Their bright red and black markings provide a striking contrast against green foliage (Plate 3*a*). Because these beetles are active during daylight hours, they are highly visible. The most familiar beetles are 6–8 mm long, oval, flat underneath, and rounded on top. Females lay clusters of 10–50 yellow-orange spindle-shaped eggs (Plate 3*b*) on foliage near aphids. Larvae do not resemble adults. Their short legs protrude from their elongate bodies, which are commonly dark with brightly colored markings (Plate 3*c* and *d*). Larvae are found in aphid colonies. Although both adult and larval ladybird beetles found on strawberry and raspberry crops prefer aphids, they will, if aphids are scarce, use food from other sources, such as small caterpillars, young plant bugs, aphid honeydew, and plant nectar. Cannibalism is common when other food sources are absent. The number of aphids eaten daily is roughly related to the size of the beetle. A full-grown larva consumes about 50 aphids per day. An adult female needs to eat about 100 aphids before she is ready to lay eggs, and she will consume at least 2400 aphids during her life-span of 1–2 months. Adults chew their victims, consuming all but the toughest parts of the body shell. Larvae usually chew a hole in the body wall and suck out the body fluids. The most common aphid-feeding species are also the most prolific, each producing several hundred eggs. Ladybird beetle adults are highly mobile and are known to converge on areas where aphids have reached high densities. In these instances, and when the weather is warm, they can be very effective in reducing aphid numbers. When an aphid population falls, however, many nonflying larvae starve because they cannot leave the area. They literally run out of food.

Development from an egg to an adult ladybird beetle requires 20–35 days, and, in temperate climates, two or three generations per year are common. Adults winter in sheltered spots, usually with many other ladybird beetles. It is not uncommon, in early fall, to see large groups of adult ladybird beetles congregating on the ground or on a building, as they prepare to search for a suitable wintering site. Every region of Canada has

several species of ladybird beetle. All are considered very effective natural predators of aphids.

Spider mite destroyers

Other coccinellid beetles, *Stethorus* species, are voracious predators of all stages of spider mites. The adult is 1–1.5 mm long, hemispherical, and glossy black (Plate 3*e*). It is not as easily observed as the highly visible red ladybird beetles but can be readily seen with the naked eye as it searches for mites on strawberry and raspberry leaves. The larvae are brown, elongate, and about 2 mm in length when mature and are sparsely covered by tufts of hair (Plate 3*f*). The pupa is a black, rough, globular form about 1 mm long and attached at one end to the surface of the leaf. The adult beetles have chewing mouthparts and generally consume the entire prey, whereas the larvae suck out the contents of the prey leaving behind an empty shell. Adults may consume 20–60 spider mites and lay up to 10 eggs per day over 1.5–3 months. Developmental time from egg to adult is from 2 to 4 weeks depending on temperature. Consumption of spider mites is slow when the larvae are small but may be as high as 250 per day in the final stages of larval growth. Various species of this group occur across Canada.

Rove beetles (Coleoptera: Staphilinidae)

Adult rove beetles are black or brown and vary in size from 1.5 to 25 mm. The body shape is characteristic of all rove beetles regardless of their size: long and slender with short wing covers (Fig. 3). The wings are folded under these short covers and a considerable length of abdomen extends behind them. Rove beetles have the habit of curving their abdomen upwards when disturbed, as do scorpions. The head is typically carried slightly bent down under the body. The beetles are very active and readily run or fly. Females lay their eggs in the soil or in decaying plant matter. Larvae are somewhat similar in appearance to adults. Many members of this group prey on other insects, especially on eggs and larvae in the soil as well as eggs and larvae of pests feeding on plants. A few small species feed on spider mites. Both adults and larvae are active and aggressive predators. Rove beetles are a large, transcontinental family with more than 800 species known from Canada.

One group of rove beetles, *Tachyporus*, with species occurring across Canada, is known to attack preferentially aphids inhabiting low-growing plants such as strawberry. In the daytime the beetles remain in the soil, but at night they leave the soil and climb the plant in search of aphids.



Plate 3 *a* **Ladybird beetle adult. (Photo by C. Chan)**
b **Ladybird beetle eggs. (Photo by C. Chan)**
c **Ladybird beetle larva. (Photo by C. Chan)**
d **Ladybird beetle larva. (Photo by C. Chan)**
e **Spider mite destroyer adult. (Photo by C. Chan)**
f **Spider mite destroyer larva. (Photo by C. Chan)**

Ground beetles (Coleoptera: Carabidae)

Ground beetles (Figs. 4–6) come in a variety of shapes, sizes, and colors. Most species are metallic blue, black, or brown, are somewhat flattened, and have faint longitudinal grooves running up and down their long wing covers. They range in size from a few millimetres to a few centimetres. Like rove beetles, most species hide by day either in the soil, under plant debris, or in other sheltered spots and are active at night. When disturbed, they run quickly but rarely fly. Nearly all species are predaceous and beneficial. Larvae are elongate and paler in color, and some have large pincer-like jaws carried on a large head. They are found in the same habitats as adults and are also predaceous. Development from egg to adult often takes 1 year and some of the larger species live for several years. The largest and usually most brightly colored ground beetles are predators of caterpillars and, on occasion, of adult weevils, whereas smaller species attack eggs, small caterpillars, and other pest insects. One large species common across Canada (*Carabus nemoralis* Müller) prefers slug and snail prey. More than 800 species are recorded from Canada.

Beneficial true bugs

Damsel bugs (Hemiptera: Nabidae)

Damsel bugs are long, slender, pale brown insects up to 10 mm long (Fig. 7). Their legs are long and the front pair is adapted for grasping prey. These predaceous bugs generally frequent low-growing plants, such as grasses and strawberry crops, where they feed on a variety of soft-bodied insects occurring on the foliage—especially aphids, leafhoppers, and some small caterpillars and moth eggs. They are swift and aggressive predators and can even capture larger lygus bug nymphs. Damsel bug nymphs resemble adults in shape but they are smaller and often lighter in color (Fig. 8). They too are predaceous.

In laboratory studies, damsel bug adults have been able to eat four lygus bug nymphs per day. Once a victim is pierced with the sharp mouthparts, it is paralyzed and never recovers, even if it is not eaten. The long, piercing mouth parts are held close to the underside of the body when they are not in use. Damsel bugs appear similar to assassin bugs and are common predators across Canada. They have been studied lately because of their potential role as biological control agents.

Assassin bugs (Hemiptera: Reduviidae)

This family is almost entirely predaceous, attacking a wide variety of plant-feeding insects, especially aphids, leafhoppers, and caterpillars. Occasionally they will attack a beneficial insect such as a ladybird beetle larva. Assassin bugs are usually longer than 10 mm, with narrow bodies and long legs (Fig. 9). They may not be distinguished easily from damsel bugs except that they are commonly larger and darker in color. Assassin



Fig. 3 Rove beetle adult. (Photo by C. Chan)

Figs. 4–6 Ground beetle adults. (Photos by H. Severson)

7



8



9

Fig. 7 Damsel bug adult. (Photo by B.D. Schaber)

Fig. 8 Damsel bug nymph. (Photo by B.D. Schaber)

Fig. 9 Assassin bug adult. (Photo by C. Chan)

bugs are usually black or brown as adults, but nymphs can be brightly colored. All stages are predaceous. Eggs are usually laid in the soil. There is commonly one generation in a year, but some species require several years to complete a single generation. Different species may pass the winter as eggs, nymphs, or adults. They are common in every part of Canada.

Assassin bugs attack by stabbing their long, somewhat curved mouthparts into a host. When they are at rest the mouthparts are not held so close to the underside of the body as the mouthparts of damsel bugs. Some species are able to paralyze their prey.

Bigeyed bugs (Hemiptera: Lygaeidae)

Adult bigeyed bugs (*Geocoris* species) measure about 4 mm long and 2 mm wide, are pinkish gray to black, and have large distinctive eyes protruding from either side of their head (Plate 4*a*). They have different color phases, which means that the pattern on the body may vary between members of the same species. Eggs are pinkish white and laid on the undersides of leaves. Nymphs resemble adults except that they are smaller. Both adults and nymphs drop quickly from the plant when disturbed. They feed on all stages of spider mite, aphid, and leafhopper. They also feed on the eggs and young nymphs of the lygus or tarnished plant bug (*Lygus lineolaris* (Palisot de Beauvois)), each consuming two to four per day. The most common species, large bigeyed bug (*Geocoris bullatus* Say) is found across Canada.

Bigeyed bugs appear in fields in the spring when lygus bugs begin to reach economically damaging levels. They are often confused with lygus bugs, especially the nymphal stages. Although their body shapes are quite similar, the large-eyed, light-brown bigeyed bugs are easy to distinguish from the small-eyed, green-bodied lygus nymphs. In addition, both adults and nymphs of the bigeyed bugs drop quickly from the plant when disturbed. Lygus nymphs and adults usually run when disturbed. Bigeyed bug nymphs and adults possess piercing and sucking mouthparts with which they attack lygus nymphs and other soft-bodied prey. In recent years, these predators have been gaining some resistance to chemicals used in lygus control (especially in areas where lygus attacks alfalfa), and so they are becoming more common and useful.

Minute pirate bugs (Hemiptera: Anthocoridae)

Adult minute pirate bugs are small, only 2 mm long, somewhat flattened, and oval in shape, with black and white wings (Plate 4*b*). Their eggs are inconspicuous because they are partially inserted into plant tissue of stems, mid ribs, or leaf petioles. Their mouthparts form long, pointed beaks with which they pierce and impale their prey. The nymphs, which are similar in shape to adults, are pale yellowish pink to brown and very active (Plate 4c). Both nymphs and adults attack and feed on all stages of spider mites as well as on thrips, aphids, and small caterpillars. Nymphs consume



Plate 4 *a* **Bigeyed bug adult. (Photo by C. Chan)**
b **Minute pirate bug adult (*Orius* sp.). (Photo by M. Herbut)**
c **Minute pirate bug nymph (*Orius* sp.). (Photo by M. Herbut)**
d **Green lacewing eggs. (Photo by R. Smith)**
e **Green lacewing larva. (Photo by C. Chan)**
f **Brown lacewing adult. (Photo by W. MacDiarmid)**

about 33 spider mites per day. Because they overwinter as adults in sheltered spots, they appear quite early in the spring and three or four generations per season can be completed outdoors. Minute pirate bugs are among the commonest predators found on many crops, including strawberry and raspberry, everywhere in Canada. They are available commercially in Canada and are used, at present, to control thrips in greenhouse crops.

Beneficial lacewings

Green lacewings (Neuroptera: Chrysopidae)

There are 25 species of green lacewings in Canada, and the adults are common and easily recognized predators of soft-bodied insect pests. The striking 14 mm long, green adult (*Chrysopa*), with its large membranous wings and commonly golden eyes, is active at night (*cover photo*). They are attracted to lights and can often be found at lighted windows or entering houses. The female lays distinctive elongate white eggs, each on a long slender stalk (10 mm), either singly or in small groups, on foliage (Plate 4*d*). The brownish larvae, which hatch from these eggs and climb down the stalks, resemble small, bristled alligators with large, curved, pincer-like jaws, which they use to grasp their prey before sucking out its body fluids (Plate 4*e*). Because of their voracious appetite for aphids, these larvae are commonly called "aphid lions." Larvae are more important predators than adults, consuming from 20 to a few hundred aphids per day. Although they prefer to feed on aphids and mealy bugs, they will attack spider mites, mite eggs, leafhoppers, small caterpillars, and thrips. Adults of some species prefer plant nectar and the honeydew excreted by aphids or mealy bugs, but many feed on insects as well.

The larvae (Plate 4*e*) reach a length of about 6 mm before pupating in a spherical, tightly woven, white cocoon on the foliage. Development from egg to adult requires 30–40 days. Adults live 4–6 weeks and a female can produce between 100 and 200 eggs. Green lacewings are among the first predators active in the cool spring weather and therefore are very important in suppressing aphid numbers early in the year. Twenty-five species are known from across Canada. Rearing methods are being developed for green lacewings so that they may be used in biological control of aphids. It is possible that they will be commercially available in the near future.

Brown lacewings (Neuroptera: Hemerobiidae)

Brown lacewing adults appear similar to green lacewings except that they are brown to gray-brown and somewhat smaller (Plate 4*f*). They are nocturnal in habit and, like the green lacewings, are often found entering lighted buildings on summer nights. Larvae are similar in shape and color to green lacewing larvae but their mouthparts are shorter and stouter. Females lay elongate eggs without stalks. Eggs are white when laid but

become orange to pink then darker brown just before hatching. In contrast to the spherical, tightly woven cocoon of green lacewings, the loosely woven cocoon is elliptical in shape and found in protected places rather than exposed on foliage.

All brown lacewing adults and larvae are predaceous on small, soft-bodied insects such as aphids, and on all stages of spider mite. Female brown lacewings lay more eggs than green lacewings and their larvae are even more common than larvae of green lacewings in some areas early in the growing season. To complete the cycle from egg to adult takes 24–30 days. Twenty three species of brown lacewings occur across Canada.

Beneficial mites

Predator mites (Acari: Phytoseiidae)

Predator mites are rarely longer than 0.5 mm and are difficult to see with the naked eye. A hand lens may be used, however, to observe predator mites on strawberry or raspberry leaves infested with spider mites. Predator mites are commonly globular or pear-shaped (Plate 5*a*), and they move more quickly than plant-feeding mites. This relative difference in speed is a good way to recognize a predator mite, but many species occur and accurate identification requires a trained eye and sophisticated microscopes. *Amblyseius fallacis* (Garman) (Plate 5*a*) is a common predator, which, with the exception of the Prairie Provinces, occurs on strawberry and raspberry crops across Canada.

Predator mites have mouthparts that they use to sense, grasp, pierce, and suck out the body fluids of prey such as plant-feeding mites, insects, and nematodes. Many predator mites also feed on plant sap, honeydew (of aphids), fungi, and pollen, which is important for maintaining the predator population when prey are not available. A female *A. fallacis* can produce one to three eggs per day, whereas other species produce an average of less than one to more than four eggs per day, depending on temperature and food availability. Development of *A. fallacis* from egg to adult requires 5–16 days. At summer temperatures (27°C), the average time for many predator species is about 6 days. Predator mites are often effective biological control agents because they develop faster than the pest species. Predator mites are available commercially in Canada and are used for control of twospotted spider mite (*Tetranychus urticae* Koch).

Other predators

Additional predators are present in any normal crop situation, often in significant numbers, but are generally not specific to the pest species present. These include both web-spinning spiders (Plate 5*b*) such as the common garden orb-spinner, and ambushing spiders such as the crab spiders. Many wasps are included in this group, such as the yellow jacket wasp (Plate 5*c*) and the baldfaced hornet (Plate 5*d*). Robber flies, which attack flying insects and capture them on the wing are also important.



Plate 5 ***a*** **Predator mite with eggs. (Photo by D.A. Raworth)**
b **Spider with prey. (Photo by W.G. Wellington)**
c **Yellow jacket wasp with caterpillar. (Photo by W.G. Wellington)**
d **Baldfaced hornet. (Photo by W.G. Wellington)**
e **Dragonfly adult. (Photo by W.G. Wellington)**
f **Lygus bug adult (*with wings*) and nymph (*without wings*). (Photo by W. MacDiarmid)**

Their larvae, which are also predaceous, attack beetle larvae (white grubs) or pupae nearer the ground. In some situations, dragonflies (Plate 5*e*) and damselflies are numerous, and both are active hunters of insect prey.

Insect-eating birds are also important in reducing numbers of insects, not just flying insects but also caterpillars and aphids. Some small rodents like moles are opportunistic feeders that will attack any large insect or pupa that they find on or in the ground.

Common insect pests of raspberry and strawberry crops

Lygus bug and other plant bugs (Hemiptera: Miridae)

Plant bugs include the lygus bug (tarnished plant bug) (Plate 5*f*), stink bug, cinch bug, and others. Most plant bugs overwinter as eggs, but a few hibernate as adults under debris in the field. In the spring, eggs hatch or adults become active and lay eggs. Lygus nymphs resemble large aphids (Plate 5*f*)—delicate, usually green, and without fully developed wings. Nymphs run quickly when disturbed. They go through five stages or molts in about 5 weeks and feed on plant juices during each stage, as well as when they are adults. Feeding on the flower blossoms causes uneven pollination, which results in "cat facing" in strawberries and "crumbly berry" in raspberries. Adults have characteristic markings on their backs and carry their long, pointed, sucking mouthparts folded under their bodies. When disturbed, they fly or run and hide.

Not all plant bugs are harmful; some are predators. Of the five bug species found on strawberry crops, all feed on plants but one will also feed on other insects. Of the 22 species found on raspberry crops in Canada, 6 are plant feeders, 10 are predators, and 6 will feed both on plants and on insect prey.

Lygus bugs are parasitized by a number of braconid wasps, at least one ichneumonid wasp, and several tachinid flies. Many insects have been recorded as predators of lygus bugs especially damsel bugs, bigeyed bugs, other plant bugs, and lacewings but little is known about how important they are. In general, these predators prefer soft-bodied prey but will also feed on lygus bugs, especially the young nymphs.

Aphids (Homoptera: Aphididae)

Aphids are small, soft-bodied, plant sap-sucking insects, usually green, black, yellowish, or pink. During most of the growing season, they deposit live young on the plant as they feed. This results in clusters or colonies of aphids of all ages (Plate 6*a*). When aphids feed, they take in large volumes of plant fluid, which they must excrete rapidly or they would explode. The liquid excreted is called honeydew because it contains a large amount of plant sugars. This shiny, sticky substance can be seen contaminating plants around aphid colonies. Large quantities of honeydew will support the growth of black sooty mold, which further contaminates the plants.

Honeydew also attracts predators and parasites of aphids for two reasons: first, many of them find aphids by first finding honeydew, and second, the sugars and amino acids in honeydew are a source of food (i.e., for adult lacewings, parasitic wasps, adult syrphid flies, aphid midges, and ladybird beetles).

The raspberry aphid is relatively large, 3 mm in length, dark green to yellow and feeds most often on the new terminal growth. The strawberry aphid (Plate 6*a*) is small and pale green, with many hairs. It overwinters as adults or as small, black eggs on older leaves. Both these aphids are important because they transmit viral diseases to their host plant.

Aphids are the favored food for a number of predatory insects including syrphids, aphid midges, lacewings, ladybird beetles, and some rove beetles. They are among the insects used by many other predaceous insects such as damsel bugs, assassin bugs, and ground beetles. They are attacked by many species of parasitic wasp especially the braconid wasps.

Leafhoppers and spittlebugs (Homoptera)

Leafhoppers, which attack raspberry, and spittlebugs, which can be a problem on strawberry, are related to aphids and damage plants in much the same way, by sucking plant juices. Leafhoppers are 3 mm long, active, slender insects, which usually jump when disturbed. They vary in color from pale white to brownish green (Plate 6*b*) and feed on the underside of young raspberry leaves. Their feeding causes yellow areas along the veins and at the edges of the leaves, giving the leaf a mottled appearance. Some will also feed on berries. Leafhoppers appear in late May in some regions, and in June and July in others. Like aphids, they excrete honeydew, which contaminates the plant or berry with a shiny, sticky substance. Black sooty mold can develop on this substance. Also like aphids, they are attacked by a variety of predators such as lacewings, ladybird beetles, some rove beetles, damsel bugs, assassin bugs, and ground beetles.

Spittlebugs have a wide distribution across Canada and, by feeding on juices of strawberry plants, cause dwarfing and low yield. The insect is covered with spittle, which is a mixture of air and their excretory product. Nymphs must have the spittle to survive. The wedge-shaped species (*Philaenus leucophthalmus* Linnaeus) that feeds on strawberry is pale white to yellow as a nymph, and grayish brown as an adult. Adults reach a length of 6 mm and move short distances between plants by flight or by jumping. They lay orange eggs on the plant, which overwinter in the field, and from which nymphs hatch in the spring. There are no good natural controls for spittlebugs.

Weevils and other beetles (Coleoptera)

A number of weevils attack the roots, crown, buds, and leaves of raspberry and strawberry plants (Plate 6*c*). Most of the damage is done when larval stages feed on roots (commonly 5–15 cm below the surface) or crown. Most weevil species attacking strawberry and raspberry crops

reproduce asexually (there are no males). None are able to fly. The adults are active at night and rarely seen in the daytime. Their presence can be detected by characteristic notching around the edges of leaves or, in the case of the strawberry clipper (Plate 6*d*), clipped blossom stems. Larvae (Plate 6*d* and *e*) can be found in the soil, feeding on the roots at any time during the growing season. Damage from crown-boring larvae causes plants to wilt at the time the fruit forms. The largest weevil is the black vine weevil (*Otiorhynchus sulcatus* (Fabricius)) at 13 mm (Plate 6*e*), and the smallest is the woods weevil (*Nemocestes incomptus* (Horn)) at about half that size.

It is important to be able to distinguish adult weevils from adult beneficial ground beetles, some of which are the same size and color. The common name for weevils is snout-nosed beetles. The adult head is narrow and placed on an elongated thorax, which looks somewhat like a neck; antennae commonly bend in the middle. Their bodies are hard and dull-colored, and they move rather slowly. Predatory ground beetles, on the other hand, move quickly when startled, are more rounded in appearance, and are usually glossy. Ground beetles, too, are nocturnal and hide under debris or in burrows in the daytime.

Weevils are not attacked by as many insects as aphids, but larvae and adults can fall prey to various vertebrates (e.g., moles) as well as to larger ground beetles and rove beetles. Eggs are also eaten by a number of small ground beetles. Other predators such as earwigs have fed on various life stages of weevils in lab studies but their importance in the field is not known.

Other beetles are also pests of raspberry and strawberry crops. The raspberry fruitworm is widely distributed within Canada. Adults feed on the new leaf material and blossoms, which reduces fruit set. Larvae penetrate the fruit, causing it to drop or to be unmarketable wormy fruit. This pest is susceptible to attack by the same variety of predators to which the weevils are susceptible. Another beetle pest is the strawberry rootworm, also of wide distribution on both strawberry and raspberry crops. Adults feed at night on leaves, whereas larvae feed on roots and strawberry crowns. Both the adult and larval stages are attacked by soil-dwelling predators such as rove beetles and ground beetles.

Leafrollers, caterpillars, climbing cutworms, and borers (Lepidoptera)

This group of pests are moths (example Plate 6*f*), the immature forms of which are caterpillars that feed on various parts of the host plant: leaves, crown, roots, and canes. Different species damage strawberry and raspberry crops to varying degrees in all areas of Canada. They are a diverse group of pests, which are attacked by a diverse group of predators and parasites; tachinid flies, ichneumonid wasps, braconid wasps, chalcid wasps, ground beetles, assassin bugs, predator wasps such as yellow jacket wasps, and, to a lesser extent, rove beetles, ladybird beetles, damsel bugs, minute pirate bugs, lacewings, and other nonspecific predators such as



Plate 6 *a* Aphids on a strawberry leaf. (Photo by W. MacDiarmid)
b *Macropsis* leafhopper (from raspberry). (Photo by H. Severson)
c Weevil on strawberry leaf. (Photo by W. MacDiarmid)
d Strawberry clipper adults and larva. (Photo by N. Bostanian)
e Black vine weevil adult, larva, and pupa. (Photo by W.T. Cram)
f Raspberry looper moth (*Autographa ampla* (Walker)). (Photo by W. MacDiarmid)



Plate 7 *a* Parasitized raspberry looper (notice the black spots). (Photo by W. MacDiarmid)
b Dead parasitized raspberry looper. (Photo by W. MacDiarmid)
c Raspberry crown borer adult. (Photo by W.G. Wellington)
d Sawfly adult. (Photo by W.T. Cram)
e Twospotted spider mite adult. (Photo by C. Chan)
f Twospotted spider mite, red wintering females. (Photo by C. Chan)

birds and rodents. A caterpillar attacked by a parasitic wasp or tachinid fly, usually shows the marks left by the female when she injects an egg (Plate 7*a*). A parasitized caterpillar which has died often has a distinctive corpse, which indicates that it was parasitized (Plate 7*b*).

Although the raspberry crown borer (*Bembicia marginata* (Harris)) is truly a moth, it looks and behaves like a yellow jacket wasp (Plate 7c). This mimicry protects it from predators such as birds; they mistake it for a wasp and expect it to sting. The adult moths are active in raspberry plantings and lay eggs on the undersides of leaves during the day, from late July until mid September. From each egg hatches a small white larva, which crawls down the cane to the soil. There it forms a small blister in the bark and overwinters. In the following spring, the larvae girdle new canes before tunneling into the soft pithy centre of the cane where they feed and spend their second winter. In their second spring, the large larvae eat their way up the canes to a few centimetres above the soil where they pupate just beneath the bark. Adult moths emerge in late July.

Because adult raspberry crown borers are so similar in appearance to syrphid fly adults and yellow jacket wasps, they may not be recognized as pests. Conversely, wasps and syrphids may be taken for the crown borer moth. The body of the moth is covered with feathery scales giving it a dull, soft appearance. Syrphid flies and yellow jacket wasps are shiny by comparison.

Sawflies (Hymenoptera: Tenthredinidae)

The raspberry sawfly (*Monophadnoides geniculatus* (Hartig)) adult is a black wasp with red and yellow markings and clear wings (Plate 7*d*). It is a stout-bodied wasp, about 6 mm long, which does not have the typical narrow wasp-like waist. It appears in raspberry plantings in May or June and lays eggs on terminal leaves of lst-year canes. The eggs hatch into spiny, green larvae, which are the same color as the leaf and therefore are difficult to see. In heavy infestations, the larvae consume the leaf tissue leaving only the larger veins. When they are fully grown, 12–17 mm long, the larvae drop to the soil and form a cocoon in which to spend the winter. The sawfly eggs are attacked by a braconid parasite; the cocoon on the ground is easy prey for ground beetles and rove beetles; and the larvae are large targets for ichneumonid wasps and tachinid flies, as well as yellow jacket wasps and birds.

Mites (Acari)

Adults of the twospotted spider mite (*Tetranychus urticae* Koch) are barely visible without magnification. They are greenish yellow with two dark spots on their back (Plate 7*e*). They feed by sucking plant sap from the undersides of leaves. Infested leaves develop a white speckling, which makes them appear dull and pale. In heavy infestations, leaves turn brown and fall off. Spider mites produce a silk webbing that covers the undersides of the leaves where they feed. Females spend the winter as red-orange

adults (Plate 7*f*) on mature foliage, under trash, in leaf litter, or in the soil at the base of the plant. In the spring, they move to the new leaves to feed and lay eggs. Hot dry conditions encourage twospotted spider mite populations to increase. Dust blowing from roadways near raspberry or strawberry plantings may have the same effect on the mite population.

Cyclamen, or strawberry, mites (*Phytonemus pallidus* (Banks)) are tiny and feed hidden in the crown and unopened leaves of strawberry plants. Even with a hand lens it is difficult to see them, but their damage is easy to detect. Unopened leaves turn yellow and some fail to open; those that do open are usually leathery and distorted. Heavily infested plants are commonly stunted, and mites feeding on the blossoms cause fruit distortion. Considerable control results from the activities of predatory mites such as *Amblyseius* species.

On the west coast an eriophyid mite called the redberry mite (*Acalitus essigi* (Hassan)) can be of importance in raspberry crops. The mites are microscopic; they spend the winter in the buds and feed on the developing fruit during spring and summer. Affected druplets remain hard and sour after the fruit is ripe. Predatory mites are of some value in controlling this pest.

Many predators feed on mites, such as some midges, the small black ladybird beetles (*Stethorus* species), some minute rove beetles, bigeyed bugs, minute pirate bugs, green and brown lacewings, and predatory mites. In southern Ontario, the predatory sixspotted thrips, *Scolothrips sexmaculatus* (Pergande), attacks spider mites. This thrips is common in the United States but has not been reported widely from Canada.

More information

Borrer, D.J.; DeLong, D.M.; Triplehorn, C.A. 1976. An introduction to the study of insects. 4th edition. Holt Reinhart & Winston.

Carr, A. 1979. Rodale's color handbook of garden insects. Rodale Press, Erasmus, Pa.

Chant, D.A. 1985. External anatomy. Pages 5–9 *in* Helle, W.; Sabelis, M.W., eds. Spider mites, their biology, natural enemies and control. Vol. 1B. Elsevier, Amsterdam.

Chant, D.A.; Hansell, R.I.C. 1971. The genus *Amblyseius* (Acari: Phytoseiidae) in Canada and Alaska. Can. J. Zool. 49:703–758.

Clausen, C.P. 1940. Entomophagous insects. McGraw Hill Book Co., N.Y.

Clancy, D.W.; Pierce, H.D. 1966. Natural enemies of some Lygus bugs. J. Econ. Entomol. 59:853–858.

Danks, H.V. 1979. Canada and its insect fauna. Entomol. Soc. Can. Memoir 108.

Davidson, R.H.; Lyon, W.F. 1979. Insect pests of farm, garden, and orchard. 7th edition. John Wiley and Sons, N.Y.

Garth, G.S.; Shanks, C.H., Jr. 1978. Some factors affecting infestation of strawberry fields by the black vine weevil in western Washington. J. Econ. Entomol. 71:443–448.

Gordon, R.D. 1985. The Coccinellidae (Coleoptera) of America north of Mexico. J. N.Y. Entomol. Soc. 93:1–912.

Kelton, L.A. 1982. Plant bugs on fruit crops in Canada. Research Branch, Agric. Can. Monograph No. 24.

Lattin, J.D. 1989. Bionomics of the Nabidae. Ann. Rev. Entomol. 34:383–400.

Lim, K.P.; Stewart, R.K. 1976. Parasitism of the tarnished plant bug, *Lygus lineolaris* (Hemiptera: Miridae) by *Peristenus pallipes* and *P. pseudopallipes* (Hymenoptera: Braconidae). Can. Entomol. 108:601–608.

McMurtry, J.A.; Huffaker, C.B.; van de Vrie, M. 1970. Ecology of tetranychid mites and their natural enemies: A review. I. Tetranychid enemies: Their biological characters and the impact of spray practices. Higardia 40:331–390.

Oatman, E.R.; Badgley, M.E.; Platner, G.R. 1985. Predators of the two-spotted spider mite on strawberry. Calif. Agric. Jan./Feb.:9–12.

Putman, W.L. 1955. The bionomics of *Stethorus punctillum* Weise (Coleoptera: Coccinellidae) in Ontario. Can. Entomol. 87:9–33.

Schneider, F. 1969. Bionomics and physiology of aphidophagous syrphidae. Ann. Rev. Entomol. 14:103–124.

Shanks, C.H., Jr. 1981. Insects and mites. Pages 293–308 *in* Childers, N.F., ed. The strawberry. Horticultural Publications, Gainsville, Fla.

Tanigoshi, L.K. 1982. Advances in knowledge of the biology of the Phytoseiidae. Pages 1–22 *in* Hoy, M.A., ed. Recent advances in knowledge of the Phytoseiidae. Univ. Calif. Press Publ. 3284.

Ware, G.W. 1980. Complete guide to pest control—with and without chemicals. Thomson Publications, Fresno, Calif.

Provincial and state publications

Berry production guide for commercial growers. 1988. B.C. Ministry of Agriculture and Fisheries, British Columbia Blueberry Co-Op Association, British Columbia Cranberry Marketing Board, British Columbia Raspberry Growers Association, and Fraser Valley Strawberry Growers Association.

Hunter, C.L. 1981. Insects and diseases of cane fruits. Ministry of Agriculture and Food, Province of Ontario. Factsheet No. 81-001, Agdex 230.605.

Johansen, C.; Eves, J.; Retan, A. 1972. Beneficial predators and parasites found on Washington crops. Washington State University Extension, Extension Bulletin 640.

Raspberry pest identification. 1988. Plant Industry Branch, New Brunswick Dep. Agric. Publ. No. 020-88, Agdex No. 232.600.

Retan, A.H.; Curtis, J. 1972*a*. Insect answers—minute pirate bug. Co-operative Extension Service, College of Agriculture, Washington State University, Pullman. E.M. 3702.

Retan, A.H.; Curtis, J. 1972*b*. Insect answers—bigeyed bugs. Co-operative Extension Service, College of Agriculture, Washington State University, Pullman. E.M. 3703.

Retan, A.H.; Curtis, J. 1972*c*. Insect answers—hover or syrphid flies. Co-operative Extension Service, College of Agriculture, Washington State University, Pullman. E.M. 3704.

Strawberry pest identification. 1988. Plant Industry Branch, New Brunswick Dep. Agric. Publ. No. 010-88, Agdex No. 232.600

Agriculture Canada

Publication 1863/F

Insectes utiles et nuisibles du fraisier et du framboisier



Canada

Insectes utiles et nuisibles du fraisier et du framboisier

Rédigé à contrat par
D.E. Henderson
E.S. Cropconsult Ltd.
Vancouver (Colombie-Britannique)

en collaboration avec
D.A. Raworth
Station de recherches
Vancouver (Colombie-Britannique)

Couverture
Chrysope à la recherche de pucerons (photo, D. Gillespie)

Agriculture Canada Publication 1863/F
On peut en obtenir des exemplaires à la
Direction générale des communications
Agriculture Canada, Ottawa (Ontario) K1A 0C7

©Ministre des Approvisionnements et Services Canada 1991
N° de cat. A53-1863/1991F ISBN 0-662-96632-5
Imprimé en 1991 2 M-04:91

Production du Service aux programmes de recherches

Also available in English under the title
Beneficial insects and common pests on strawberry and raspberry crops

Table des matières

Remerciements 6

Introduction 7

Diptères utiles (mouches) 10
Syrphes (Diptères : syrphidés) 10
Tachinaires (Diptères : tachinidés) 10
Cécidomyies (Diptères : cécidomyiidés) 13

Hyménoptères utiles (guêpes) 13
Ichneumons (Hyménoptères : ichneumonidés) 13
Braconides (Hyménoptères : braconidés) 14
Chalcidies (Hyménoptères : chalcidoïdés) 15

Coléoptères utiles 15
Coccinelles (Coléoptères : coccinellidés) 15
Staphylins (Coléoptères : staphilinidés) 20
Carabes (Coléoptères : carabidés) 20

Hémiptères utiles 22
Punaises demoiselles (Hémiptères : nabidés) 22
Réduves (Hémiptères : réduviidés) 22
Punaises tête-de-clou (Hémiptères : lygéidés) 23
Anthocoris (Hémiptères : anthocoridés) 23

Névroptères utiles 25
Chrysopes (Névroptères : chrysopidés) 25
Hémérobies (Névroptères : hémérobidés) 25

Acariens utiles 26
Acariens prédateurs (Acariens : phytoséidés) 26

Autres prédateurs 26

Ravageurs courants du framboisier et du fraisier 27
Punaises et lygides (Hémiptères : miridés) 27
Pucerons (Homoptères : aphididés) 27
Cicadelles et cercopes (Homoptères) 30
Charançons et autres coléoptères (Coléoptères) 31
Enrouleuses, chenilles, noctuelles et perceurs (Lépidoptères) 32
Tenthrèdes (Hyménoptères : tenthrèdinidés) 32
Acariens 33

Bibliographie 34

Remerciements

La présente brochure a vu le jour en réponse à une demande des membres de la *Lower Mainland Horticultural Improvement Association*, formulée en février 1988, lors des mini-cours présentés chaque année aux cultivateurs, à Clearbrooke (C.-B.). Il s'agit en fait d'un projet coopératif. Le Service aux programmes de recherches d'Agriculture Canada s'est occupé de la publication, dont il a absorbé le coût, tandis que la Direction de la protection des cultures du ministère de l'Agriculture et des Pêcheries de Colombie-Britannique a financé la collecte des données et la rédaction du document en sous-traitance par des spécialistes. Enfin, les chercheurs et les professionnels des quatre coins du Canada dont voici la liste ont fourni photographies, spécimens et renseignements sur les insectes de leur région : R. Smith, Vedalia Research, Galiano (C.-B.); W.G. Wellington et S.G. Cannings, Université de la Colombie-Britannique, Vancouver (C.-B.); D. Gillespie, Agriculture Canada, Agassiz (C.-B.); C. Vincent et N. Bostanian, Agriculture Canada, Saint-Jean-sur-Richelieu (Québec); M. Luffman, Agriculture Canada, Bouctouche (N.-B.); R. Tremblay, ministère de l'Agriculture du Nouveau-Brunswick, Fredericton (N.-B.); B.D. Schaber, Agriculture Canada, Lethbridge (Alb.); L. Gilkeson, Applied Bionomics, Sidney (C.-B.); M. Steiner et M. Herbut, Centre environnemental, Vegreville (Alb.); R. Long et M. Mackauer, Université Simon Fraser, Burnaby (C.-B.); C. Chan, W. MacDiarmid, H. Severson et W.T. Cram, Agriculture Canada, Vancouver (C.-B.). Ont prêté leur concours à la correction du manuscrit : H. Gerber et W. Peters, ministère de l'Agriculture et des Pêcheries de la Colombie-Britannique, Abbotsford (C.-B.) et S. Krannitz, Pherotech Ltd., Vancouver (C.-B.). Nos plus sincères remerciements à tous.

Introduction

Par crainte de détériorer l'environnement, beaucoup d'agriculteurs freinent leur usage de produits chimiques et cherchent d'autres moyens pour lutter contre les ravageurs et les maladies des cultures. Cette approche, baptisée lutte intégrée, gagne de plus en plus de popularité chaque année au Canada et ailleurs. Avant de l'adopter cependant, il faut connaître les autres méthodes de lutte et savoir comment les mettre en pratique. Ce manuel devrait fournir au moins quelques indications à cet égard. Les insectes utiles qui s'attaquent aux ravageurs sont très sensibles aux pesticides, si bien que peu d'entre eux survivent dans les champs où l'on pulvérise régulièrement des antiparasitaires. C'est aussi pourquoi beaucoup de cultivateurs, et même de vulgarisateurs, les connaissent mal. Un agriculteur qui opte pour la lutte intégrée, de son propre chef ou sur les conseils d'un expert, doit commencer par observer régulièrement la population de ravageurs afin d'évaluer la menace qu'ils posent. Il ne recourra à un insecticide que si les parasites menacent d'augmenter au point de causer des pertes économiques. Alors, la population de nombreux insectes, y compris d'insectes utiles, commencera à s'accroître dans le champ. Il est donc capital de pouvoir identifier les insectes utiles pour les différencier des ravageurs. Même s'il existe plusieurs brochures ou publications sur les ravageurs au Canada, aucune ne porte sur les insectes utiles.

La présente brochure décrit les insectes utiles du fraisier et du framboisier, cependant on retrouvera beaucoup de ces insectes sur d'autres cultures attaquées par les mêmes ravageurs. Ainsi, les prédateurs des pucerons du fraisier détruiront les pucerons, peu importe la plante sur laquelle ils se trouvent. Par ailleurs, si certains insectes utiles (les guêpes parasites par exemple) ne s'attaquent qu'à un seul ravageur, d'autres insectes ayant une forte ressemblance à ceux décrits ici seront les prédateurs d'autres espèces nuisibles. Cette brochure, par conséquent, facilitera non seulement l'identification des insectes utiles du fraisier et du framboisier, mais aussi celle des insectes utiles de la majorité des cultures, en général.



Planche 1

a **Syrphe adulte (photo W.G. Wellington)**
b **Syrphe adulte (photo W.G. Wellington)**
c **Larve de syrphe (photo C. Chan)**
d **Larve de syrphe (photo H. Severson)**
e **Larve de cécidomyie (photo W. McDiarmid)**
f **Cécidomyie attaquant un puceron (photo L. Gilkeson)**

Appel aux professionnel Quand nous avons rassemblé l'information et la documentation nécessaires à la rédaction de cet ouvrage, nous nous sommes aperçus que les renseignements manquaient sur la distribution des insectes utiles et que les photos étaient rares. Il est donc possible que les données d'ordre biologique soient incomplètes et que certaines illustrations fassent défaut. Si vous possédez des renseignements sur les insectes utiles de votre région et surtout de bonnes photographies, nous vous serions reconnaissants de communiquer avec nous. Nous avons l'intention de préparer d'autres guides du même genre pour différentes cultures. Toute contribution, sous forme d'information ou de photographies, ne pourra qu'en accroître l'utilité. Il est possible de communiquer avec l'un ou l'autre auteur aux adresses suivantes :

D.E. Henderson, E.S Cropconsult Ltd., 2962 West 15th Ave., Vancouver, B.C. V6K 3A3

D.A. Raworth, Research Station, Research Branch, Agriculture Canada, 6660 N.W. Marine Drive, Vancouver, B.C. V6T 1X2.

Diptères utiles (mouches)

Syrphes (Diptères : syrphidés)

Les syrphes sont des mouches de 6 à 13 mm de longueur. Leur abdomen rayé ressemble à celui des guêpes et des abeilles, mais il est dépourvu de dard (planche 1*a* et *b*). Les rayures servent d'avertissement et mettent l'insecte à l'abri d'éventuels prédateurs, par mimétisme. Les nombreuses espèces que compte cette famille sont de bons pollinisateurs, et leur efficacité à ce titre n'est surpassée que par les abeilles. Les syrphes se nourrissent de pollen, de nectar et de miellat (la substance collante et luisante qui macule les plantes près des colonies de pucerons). Leur aptitude à voler sur place, comme de petits hélicoptères, est ce qui les distingue le mieux des abeilles. Les syrphes ou leur larve (le stade immature) se nourrissent exclusivement de pucerons.

La mouche dépose la plupart de ses œufs sur les plantes fortement infestées, parmi les grosses colonies de pucerons. La femelle pond entre 400 et 1 000 œufs blancs oblongs qu'elle dépose un à un ou en petits amas, selon l'espèce. Quelques jours plus tard, les œufs éclosent et une minuscule larve semblable à une limace en sort. En se développant, la larve prend la couleur caractéristique de l'espèce. Certaines sont vertes, lignées ou tachées de blanc (planche 1*c*) tandis que d'autres sont jaunâtres ou brunâtres avec des marbrures (planche 1*d*). La larve cherche les pucerons en soulevant le devant du corps et en se balançant latéralement et de haut en bas, pour fouiller la surface des feuilles. Quand elle entre en contact avec un puceron, elle le perce de ses pièces buccales acérées, le soulève et le vide complètement de son contenu. Une larve peut dévorer à elle seule de 200 à 800 pucerons durant sa courte vie de 7 à 10 jours. Parvenue à maturité (elle mesure alors 6 mm), la larve se met en quête d'un endroit sombre, une feuille morte enroulée sur la plante par exemple, ou s'enfouit dans le sol. Alors commence la pupaison (stade de repos), qui durera environ 1 semaine. La jeune mouche adulte se mettra à la recherche d'un compagnon dès son émergence et le cycle recommencera. Les syrphes comptent plusieurs générations par année et sont considérés comme un prédateur très efficace des pucerons.

Tachinaires (Diptères : tachinidés)

Les tachinaires, qui regroupent de nombreuses espèces, ressemblent à la mouche domestique, à l'abeille ou à la guêpe, dont ils ont à peu près la même taille (fig. 1). On les considère comme faisant partie des insectes parasites les plus communs. L'adulte se nourrit des sucs végétaux ainsi que du miellat sécrété par les pucerons, les kermès et les cicadelles. La larve, par contre, parasite d'autres insectes, principalement les formes immatures des papillons nocturnes, des tenthrèdes et des coléoptères, mais aussi quelques punaises et des criquets. Certaines espèces n'attaquent que les chenilles qui vivent cachées, comme les enrouleuses, alors que d'autres



Fig. 1 Tachinaire parasite de l'arpenteuse de framboisier (photo W.T. Cram)



Fig. 2 Formes courantes des ichneumons, des braconides et des chalcidies



Planche 2

a **Ichneumon (*Apophua*) sur le cocon d'une enrouleuse (photo D. Gillespie)**
b **Braconide injectant son œuf (photo R. Long)**
c **«Momies» dans une colonie de pucerons (photo M. Mackauer)**
d **Pupe d'un braconide dans une momie (photo M. Mackauer)**
e **Émergence du braconide (photo C. Chan)**
f **Cocon de braconide (*Macrocentrus*) (photo D. Gillespie)**

préfèrent les noctuelles. Plusieurs espèces ont été introduites d'Europe pour lutter contre les ravageurs d'Amérique du Nord. C'est notamment le cas du tachinaire de la noctuelle, ennemi commun de ce papillon dans de nombreuses régions du Canada. La femelle dépose ses œufs directement sur l'hôte ou le feuillage qu'il dévorera. Parfois, elle les lui inocule même directement dans le corps. Quelques espèces sont vivipares et laissent les jeunes larves là où d'autres tachinaires déposeraient leurs œufs. Des œufs blancs sur une chenille indiquent que celle-ci est parasitée par un tachinaire. La larve nouvellement éclose se creuse un passage dans l'hôte dont elle dévorera les organes internes plusieurs jours durant, puis abandonnera la dépouille pour entamer la pupaison, à proximité. Tous les insectes parasités par les tachinaires finissent par mourir.

Cécidomyies (Diptères : cécidomyiidés)

Adulte, la cécidomyie du puceron (*Aphidoletes aphidimyza* (Rondani)) mesure 2 ou 3 mm de longueur. Elle est difficile à repérer en raison de ses habitudes nocturnes et de la taille minuscule de son corps délicat (planche 1*e*). La femelle vit environ 10 jours et pond près de 250 œufs. Un repas de sève ou de nectar lui permettra de produire un premier lot d'environ 40 œufs. Pour pondre davantage cependant, elle a besoin de miellat de puceron. Contrairement à l'adulte, d'aspect fragile, la petite larve orange (1,5 mm de long) est très robuste (planche 1*f*). Elle se nourrit de pucerons et reste à l'extérieur tout l'hiver, dans la plupart des régions du Canada. La larve capture des pucerons en perçant une patte et en inoculant une toxine paralysante; ensuite, elle vide complètement le corps de son contenu. Si les pucerons sont nombreux, la larve en tue plus qu'elle en mange. Après 1 ou 2 semaines de ce manège, elle se laisse choir sur le sol où elle creuse une galerie et commence la pupaison. Dix à 14 jours plus tard émerge un nouvel adulte. Des expériences effectuées en Colombie-Britannique indiquent que la population de cécidomyies du puceron augmente quand les pucerons prolifèrent et diminue lorsqu'ils se font plus rares. On peut acheter au Canada une espèce de cécidomyie qu'on utilise couramment pour combattre les pucerons dans les cultures de serre et les arbres fruitiers. Certains membres de cette famille détruisent aussi les œufs et les larves d'acariens. Enfin, d'autres parasitent les cochenilles et leurs œufs.

Hyménoptères utiles (guêpes)

Ichneumons (Hyménoptères : ichneumonidés)

Les ichneumons sont des guêpes au corps élancé qui peuvent mesurer de 5 à 36 mm de longueur. La femelle porte habituellement au bout de l'abdomen une longue structure similaire à une aiguille (tarière) qu'elle utilise pour déposer ses œufs (fig. 2, p. 11). La tarière est plus longue que le corps et on la prend souvent pour un dard, bien que son rôle consiste à introduire des œufs dans le corps des insectes appelés à nourrir la larve. La larve des ichneumons parasite les stades immatures d'autres insectes,

surtout des chenilles de papillons nocturnes ou diurnes. La longueur de la tarière permet à la femelle d'inoculer ses œufs dans des hôtes invisibles comme les enrouleuses ou les perce-tiges. Les espèces du genre *Apophua* en sont un parfait exemple puisqu'elles s'attaquent à la tordeuse du fraisier (*Ancylis comptana* (Froelich)) alors que celle-ci se cache à l'intérieur de la feuille (planche 2*a*, p. 12). Beaucoup d'ichneumons sont relativement courants au Canada. L'adulte peut être brun, rouge ou noir et porter différentes marques. Il se distingue des autres guêpes parasites par sa grande taille et la longueur de son abdomen, qui dépasse celle de la tête et du thorax réunis. Il possède également une longue tarière.

Braconides (Hyménoptères : braconidés)

Les braconides sont plus petits que les ichneumons (soit jusqu'à 12 mm de longueur). Leur corps plus court est également plus trapu (fig. 2). Les espèces de cette famille possèdent également une tarière, mais celle-ci reste cachée dans le corps dont elle ne sort qu'au moment de l'inoculation. Les braconides parasitent une plus vaste gamme d'insectes que les ichneumons, les hôtes allant de diverses chenilles comme la tordeuse des citrus (*Argyrotaeni citrana* (Fernald)) à des mouches, des guêpes, des coléoptères et des pucerons. Lorsque la femelle a injecté son œuf, il sort de celui-ci une larve qui dévorera lentement l'hôte. La larve parvient à maturité quand meurt ce dernier. Elle entame sa pupaison à l'intérieur de la dépouille ou près de celle-ci, quelque fois dans un cocon soyeux d'où émergera la guêpe adulte. À l'instar des ichneumons, on trouve de nombreuses espèces de braconides un peu partout au Canada.

Les braconides parasites des pucerons constituent un groupe important d'insectes utiles du fraisier et du framboisier. En effet, les parasites des pucerons comptent beaucoup de braconides ayant un cycle biologique analogue. La femelle inocule son œuf dans un jeune puceron avec la rapidité de l'éclair (planche 2*b*) pendant que celui-ci pompe la sève de la plante. En règle générale, l'attaque dérange le puceron dans son travail sans toutefois l'inciter à quitter la plante sur laquelle il se trouve. Le puceron parasité ne se reproduit habituellement pas, mais continue de se nourrir jusqu'à ce qu'il ait été entièrement dévoré de l'intérieur. La larve adulte fixe la carcasse du puceron à la surface de la feuille et la dépouille prend un aspect parcheminé ou noirâtre. On appelle les pucerons morts de cette façon des «momies». Ils sont très faciles à reconnaître dans une colonie de pucerons vivants (planche 2*c*). La momie protège le braconide durant la pupaison (planche 2*d*). À la fin de cette dernière, environ une semaine plus tard, la guêpe adulte découpe un trou circulaire dans la momie et sort au grand jour (planche 2*e*). La momie vide, percée d'un trou, reste sur la feuille. La présence de momies dans une colonie indique que beaucoup de pucerons sont parasités, même s'ils paraissent en bonne santé, et mourront au bout de quelques jours.

Un autre grand groupe de braconides parasitent les chenilles. Leur cycle biologique reproduit largement celui des braconides qui s'attaquent aux pucerons. Les espèces du genre *Copidosoma* font toutefois bande à

part. En effet, la femelle adulte pond son œuf dans un œuf de l'hôte. À mesure que celui-ci traverse les différents stades de son cycle, l'œuf du braconide se divise pour donner plusieurs petites larves. À la mort de l'hôte, les larves entament la pupaison et donneront naissance à plusieurs guêpes adultes. Par cet artifice, la femelle parvient à produire une progéniture nombreuse, même si elle ne trouve qu'un hôte. D'autres braconides, les espèces du genre *Macrocentrus* par exemple, tirent parti d'un hôte de grande taille comme une chenille en lui injectant un grand nombre d'œufs. Les larves grandissent en dévorant la chair de la chenille. Lorsque celle-ci meurt, on trouve regroupées dans le reste de son cocon les pupes du parasite (planche 2*f*).

Chalcidies (Hyménoptères : chalcidoïdés)

La plupart des guêpes de ce groupe sont encore plus petites que les ichneumons et les braconides. D'un bleu ou d'un vert métallique foncé, elles ne mesurent au maximum que 7 mm de longueur et ont des ailes translucides. La guêpe paraît plutôt sauter que prendre son envol. On trouve les chalcidies partout, mais on a tendance à ne pas les remarquer en raison de leur taille minuscule. Les membres de cette famille parasitent les chenilles des papillons nocturnes et diurnes, les larves de certaines mouches et celles de quelques coléoptères. Les chalcidies ne sont pas toutes utiles, car quelques-unes s'attaquent aux tachinaires et aux pupes des ichneumons.

Un groupe bien connu, le genre *Trichogramma*, pond ses œufs dans ceux de nombreuses espèces de papillons nocturnes et diurnes. L'adulte, qui mesure moins de 1 mm de longueur, montre une grande efficacité comme agent de lutte biologique. Il est possible de s'en procurer dans le commerce et on s'en sert sur diverses cultures pour détruire les chenilles, quoique moins fréquemment au Canada qu'aux États-Unis.

Coléoptères utiles

Coccinelles (Coléoptères : coccinellidés)

Coccinelles

La coccinelle adulte est sans doute le prédateur le plus facile à identifier dans un champ. Ses taches rouge vif et noires ressortent de façon frappante sur le vert du feuillage (planche 3*a*, p.16). En outre, comme elle est active le jour, la coccinelle se repère aisément. L'espèce la plus familière, de forme ovale, mesure de 6 à 8 mm de longueur, a le ventre plat et le dos arrondi. La femelle dépose un amas de 10 à 50 œufs jaune orange en forme de fuseau (planche 3*b*) sur les feuilles, près des pucerons. La larve ne ressemble pas du tout à l'adulte. Ses courtes pattes dépassent du corps allongé, souvent foncé, mais agrémenté de marques vives (planches 3*c* et *d*). La larve passe sa vie dans les colonies de pucerons. Quoi qu'elles préfèrent les pucerons, la coccinelle adulte qu'on trouve sur le fraisier et le framboisier et sa larve



Planche 3

a **Coccinelle adulte (photo C. Chan)**
b **Oeuf de coccinelle (photo C. Chan)**
c **Larve de coccinelle (photo C. Chan)**
d **Larve de coccinelle (photo C. Chan)**
e **Stethorus adulte (photo C. Chan)**
f **Larve de stethorus (photo C. Chan)**

chercheront d'autre nourriture si les pucerons se font rares, par exemple de petites chenilles, de jeunes punaises, du miellat ou du nectar. Faute d'autres sources d'aliments, la coccinelle est souvent atteinte de cannibalisme. Le nombre de pucerons dévorés chaque jour dépend approximativement de la taille du coléoptère. À maturité, la larve en détruit près de 50 par jour. La femelle adulte doit en manger environ une centaine avant de commencer à pondre et elle en dévorera au moins 2 400 durant sa vie qui dure 1 ou 2 mois. L'adulte broie sa victime et ne laisse que les parties les plus coriaces de la carapace. Habituellement, la larve perfore cette dernière et aspire les liquides du corps. L'espèce la plus commune à se nourrir de pucerons est aussi la plus prolifique, car elle pond plusieurs centaines d'œufs à la fois. La coccinelle adulte se déplace beaucoup et on en a vu converger vers les endroits où les pucerons ont atteint une grande densité. Dans ce cas, elle s'avère particulièrement efficace, surtout par temps chaud. Toutefois, quand le nombre de pucerons diminue, les larves, incapables de voler, périssent faute de pouvoir gagner un autre endroit. Elles meurent littéralement de faim.

Le cycle biologique dure de 20 à 35 jours et, en climat tempéré, il n'est pas rare de compter deux ou trois générations par année. L'adulte traverse l'hiver dans des endroits abrités, souvent en compagnie de beaucoup de ses congénères. À l'automne, il est fréquent de voir un grand nombre de coccinelles adultes se rassembler sur le sol ou sur un bâtiment en quête d'un endroit pour hiverner. Toutes les régions du Canada comptent plusieurs espèces de coccinelles qui sont des prédateurs naturels très efficaces du puceron.

Stethorus

D'autres coccinellidés, les espèces du genre *Stethorus* par exemple, sont des prédateurs particulièrement voraces qui s'attaquent à tous les stades du tétranyque. L'adulte mesure de 1 à 1,5 mm de longueur et son corps hémisphérique est noir luisant (planche 3*e*). Il est beaucoup moins facile à repérer que les coccinelles rouges, très évidentes, mais on l'aperçoit facilement à l'œil nu quand il se met en quête de tétranyques sur les feuilles de fraisier et de framboisier. La larve brune, allongée, mesure environ 2 mm de longueur à maturité et porte des touffes de poil ici et là (planche 3*f*). La pupe, de forme globulaire, a une carapace robuste et noire d'environ 1 mm de longueur. Elle se fixe à la surface de la feuille par une extrémité. Le coléoptère adulte est doté de pièces buccales qui lui permettent de dévorer sa proie en entier, alors que la larve se contente d'en sucer le contenu pour ne laisser derrière elle qu'une coquille vide. L'adulte peut détruire de 20 à 60 tétranyques par jour et pondre quotidiennement jusqu'à 10 œufs pendant 1,5 à 3 mois. Il faut de 2 à 4 semaines selon la température pour que l'œuf donne un adulte. Au début de sa vie, la larve consomme moins de tétranyques, mais vers la fin, elle peut en dévorer jusqu'à 250 par jour. Diverses espèces de ce groupe existent un peu partout au Canada.

3

4

5

6

Fig. 3 Staphylin adulte (photo C. Chan)

Fig. 4–6 Carabe adulte (photo H. Severson)

7

Fig. 7 Odonate adulte (photo B.D. Schaber)

8

Fig. 8 Nymphe d'odonate (photo B.D. Schaber)

9

Fig. 9 Réduve adulte (photo C. Chan)

Staphylins (Coléoptères : staphilinidés)

Le staphylin adulte peut être noir ou brun et sa taille varie entre 1,5 et 25 mm. Le corps a une forme caractéristique, propre aux staphylins, peu importe leur taille : longue et élancée avec de courts élytres (fig. 3, p. 18). Les ailes sont repliées sous les petits élytres et l'abdomen se prolonge considérablement derrière elles. Le staphylin a pour habitude de replier l'abdomen vers le haut quand on le dérange, comme le font les scorpions. L'insecte porte typiquement la tête légèrement repliée vers le bas, sous le corps. Il s'agit de coléoptères très actifs qui s'enfuient ou s'envolent rapidement. La femelle enfouit ses œufs dans le sol ou les débris végétaux en décomposition. La larve ressemble un peu à l'adulte. Maintes espèces de ce groupe s'attaquent à d'autres insectes, surtout les œufs et les larves dans le sol, mais elles détruisent également les œufs et les larves des phytoparasites. Un petit nombre d'espèces s'attaquent aux tétranyques. L'adulte comme la larve sont très actifs et s'avèrent des prédateurs très agressifs. Les staphylins constituent une vaste famille qu'on retrouve sur tout le continent et regroupent au-delà de 800 espèces, rien qu'au Canada.

Un groupe particulier, les espèces du genre *Tachyporus*, qu'on trouve aux quatre coins du Canada, préfère attaquer les pucerons qui habitent les plantes basses comme le fraisier. Durant le jour, le coléoptère reste sur le sol, mais la nuit venue, il grimpe dans les plantes à la recherche des pucerons.

Carabes (Coléoptères : carabidés)

Les carabes (fig. 4 à 6, p. 18) épousent de multiples formes, tailles et couleurs. La majorité des espèces sont bleues, noires ou brun métallique, ont le corps légèrement aplati et de longs élytres finement nervurés de façon longitudinale. La taille varie de quelques millimètres à quelques centimètres. Comme les staphylins, la majorité des carabes se cachent dans le sol, sous les débris végétaux ou dans des endroits similaires durant le jour et ne commencent à s'activer que la nuit. Lorsqu'ils sont dérangés, ils fuient rapidement, mais s'envolent rarement. Presque toutes les espèces sont des prédateurs utiles. La larve, allongée et plus pâle, est munie parfois de grosses mâchoires en forme de pinces au bout d'une grosse tête. On la trouve dans le même habitat que l'adulte et, comme lui, elle s'attaque à d'autres insectes. Le cycle biologique dure souvent un an et quelques espèces parmi les plus grandes vivent plusieurs années. Les plus gros carabes, souvent ceux qui ont les couleurs les plus vives, attaquent les chenilles et, à l'occasion, les charançons adultes, tandis que les espèces plus petites dévorent les œufs, les petites chenilles et d'autres ravageurs. Une grosse espèce commune au Canada (*Carabus nemoralis* Müller) préfère les limaces et les escargots. On a recensé plus de 800 espèces de carabes au Canada.



Planche 4

a **Punaise des bois adulte (photo C. Chan)**
b **Anthocoris adulte (photo M. Herbut)**
c **Nymphe d'anthocoris (photo M. Herbut)**
d **Oeufs de chrysope (photo R. Smith)**
e **Larve de chrysope (photo C. Chan)**
f **Hémérobie adulte (photo M. MacDiarmid)**

Hémiptères utiles

Punaises demoiselles (Hémiptères : nabidés)

La punaise demoiselle est un insecte au corps long, élancé et brun pâle qui peut mesurer jusqu'à 10 mm de longueur (fig. 7, p. 19). Il est doté de longues pattes dont la première paire est adaptée pour la capture des proies. Ce prédateur affectionne les plantes basses comme les graminées et le fraisier, où il se nourrit des insectes à corps mou qui vivent sur le feuillage, notamment les pucerons, les cicadelles et certaines chenilles, de même que les œufs de papillons nocturnes. Rapide et agressif, la punaise demoiselle attrape même les nymphes de lygide beaucoup plus grosses. La nymphe ressemble à l'adulte par sa forme, mais est beaucoup plus petite et de couleur plus pâle (fig. 8, p. 19). Elle a également un comportement de prédateur.

En laboratoire, on a vu un odonate adulte dévorer quatre nymphes de lygide par jour. Une fois piquée par les pièces buccales acérées, la victime reste paralysée à jamais, même si elle ne se fait pas dévorer. Au repos, les longues pièces buccales broyeuses restent plaquées sur la face ventrale du corps. La punaise demoiselle ressemble au réduve et on le retrouve couramment au Canada. Ce genre a récemment attiré l'attention des chercheurs en raison de la possibilité de son utilisation comme agent de lutte biologique.

Réduves (Hémiptères : réduviidés)

Il s'agit d'une famille d'insectes presque tous prédateurs, qui s'attaquent à une vaste gamme de phytoparasites, notamment les pucerons, les cicadelles et les chenilles. Il arrive cependant qu'ils inscrivent à leur tableau de chasse des insectes utiles comme la larve des coccinelles. Le réduve mesure habituellement plus de 10 mm de long, avec un corps étroit et des pattes filiformes (fig. 9, p. 19). Il est parfois difficile de le différencier des odonates, sauf par la taille, plus importante, et par la couleur, plus foncée. En règle générale, le réduve adulte est noir ou brun, mais la nymphe arbore parfois des couleurs vives. Tous les stades sont prédateurs. La femelle pond habituellement ses œufs dans le sol. En règle générale, on ne compte qu'une génération par année, mais certaines espèces prennent plusieurs années pour compléter leur cycle. Ainsi, quelques espèces peuvent hiverner simultanément sous forme d'œuf, de nymphe ou d'adulte. On trouve des réduves dans toutes les régions du Canada.

Le réduve attaque sa proie en la poignardant avec sa longue pièce buccale légèrement recourbée. Au repos, le réduve ne garde pas son appareil buccal aussi près du corps que l'odonate. Certaines espèces paralysent leur proie avant de la dévorer.

Punaises tête-de-clou (Hémiptères : lygéidés)

La punaise tête-de-clou adulte (espèces du genre *Geocoris*) mesure environ 4 mm de longueur et 2 mm de largeur. De couleur gris rosé à noir, elle se remarque par ses deux gros yeux qui dépassent de chaque côté de la tête (planche 4*a*, p. 21). L'espèce traverse différentes phases de couleur, ce qui signifie que le motif de la carapace peut varier entre les membres de la même espèce. Les œufs, blanc rosé, sont déposés en amas sur la face inférieure des feuilles. La nymphe ressemble à l'adulte en plus petit. L'adulte et la nymphe se laissent tomber sur le sol quand on les dérange. Ils s'attaquent à tous les stades du tétranyque, du puceron et de la cicadelle. Ils se nourrissent des œufs et des jeunes nymphes de lygide et de punaise terne (*Lygus lineolaris* (Palisot de Beauvois)), à raison de deux à quatre par jour. L'espèce la plus courante, (*Geocoris bullatus* Say), existe un peu partout au Canada.

On trouve les punaises des bois dans les champs, au printemps, lorsque la population de lygides atteint un point où elle pourrait entraîner de grandes pertes économiques. On les confond souvent avec les lygides, surtout au stade de nymphe. Pourtant si la forme du corps est similaire, la punaise des bois, avec ses grands yeux et sa couleur brun clair, se distingue aisément des nymphes de lygide, de couleur verte et aux yeux petits. En outre, l'adulte et la nymphe de punaise des bois se laissent choir sur le sol quand on les dérange tandis que la lygide s'enfuit. La nymphe et la punaise des bois adulte sont pourvues de pièces buccales qui leur permettent de percer et de sucer les nymphes de lygide et d'autres proies sans carapace. Depuis quelques années, ces prédateurs ont acquis une certaine résistance aux pesticides utilisés pour détruire les lygides (surtout dans les régions où ce ravageur s'attaque à la luzerne). Ils deviennent donc de plus en plus communs et utiles.

Anthocoris (Hémiptères : anthocoridés)

L'anthocoris est une petite punaise ne mesurant que 2 mm, au corps légèrement aplati et ovale. Ses ailes sont noir et blanc (planche 4*b*). Les œufs passent inaperçus, car ils sont partiellement enfouis dans la tige, la nervure principale ou le pétiole des feuilles. Les pièces buccales imitent un long bec pointu avec lequel l'insecte perce et empale sa proie. La nymphe, qui ressemble à l'adulte par sa forme, est de couleur rose jaunâtre pâle à brun et s'avère très active (planche 4*c*). La nymphe et l'adulte dévorent tous les stades du tétranyque de même que les thrips, les pucerons et les petites chenilles. La nymphe mange environ 33 tétranyques par jour. Comme l'adulte passe l'hiver à l'abri dans des endroits couverts, il réapparaît très tôt au printemps. Trois ou quatre générations peuvent ainsi se succéder chaque saison. Au Canada, l'anthocoris est le plus commun des prédateurs sur de nombreuses cultures, notamment le fraisier et le framboisier. On peut d'ailleurs s'en procurer dans le commerce et à l'heure actuelle, on s'en sert pour lutter contre les thrips dans les serres.



Planche 5

a **Phytoséidé et ses œufs (photo D.A. Raworth)**
b **Araignée et sa proie (photo W.G. Wellington)**
c **Guêpe jaune et chenille (photo W.G. Wellington)**
d **Guêpe jaune (photo W.G. Wellington)**
e **Libellule adulte (photo W.G. Wellington)**
f **Lygide adulte (*avec ailes*) et sa nymphe (*sans ailes*) (photo W. MacDiarmid)**

Névroptères utiles

Chrysopes (Névroptères : chrysopidés)

On compte 25 espèces de chrysopes au Canada. Ce prédateur des ravageurs à corps mou est relativement commun et facile à identifier. Long de 14 mm, l'adulte (*Chrysopa*) a un corps vert doté de grandes ailes membraneuses et d'yeux souvent dorés. Il s'active surtout la nuit (*photo de couverture*). Attiré par la lumière, il se pose fréquemment sur la vitre des fenêtres éclairées ou se faufile dans les maisons. La femelle pond des œufs caractéristiques blancs et allongés, seuls ou en petits groupes sur les feuilles, chacun à l'extrémité d'un long pédoncule de 10 mm (planche 4*d*). La larve brunâtre qui sort de l'œuf puis descend le pédoncule ressemble à un petit alligator poilu aux grosses mâchoires recourbées en forme de pinces. Ces mâchoires servent à attraper les proies et à en extraire tous les liquides organiques (planche 4*e*). En raison de sa voracité, on l'appelle couramment le lion des pucerons. La larve est un prédateur plus important que l'adulte, car elle peut dévorer d'une vingtaine à quelques centaines de pucerons par jour. Même si elle affectionne les pucerons et les cochenilles, elle ne dédaigne pas les tétranyques, les œufs d'acariens, les cicadelles, les petites chenilles et les thrips. L'adulte de certaines espèces préfère le nectar des plantes et le miellat des pucerons ou des cochenilles, mais beaucoup dévorent également des insectes.

La larve (planche 4*e*) doit mesurer environ 6 mm de long avant d'entamer la pupaison. La pupe sphérique ressemble à un cocon blanc au fil très serré, déposé sur le feuillage. Le cycle biologique complet dure de 30 à 40 jours. Durant sa vie de 4 à 6 semaines, la femelle adulte peut pondre de 100 à 200 œufs. La chrysope est le premier prédateur à se manifester au printemps, quand le temps est encore frais. Elle revêt donc une grande importance dans la lutte contre les pucerons au début de la période végétative. Au Canada, on en connaît 25 espèces. On tente actuellement de mettre au point des techniques d'élevage en vue d'utiliser cet insecte dans la lutte biologique contre les pucerons. Il est donc possible qu'on puisse s'en procurer dans le commerce dans un proche avenir.

Hémérobies (Névroptères : hémérobidés)

L'hémérobie ressemble à la chrysope sauf pour la couleur, qui va du brun au gris brun, et pour la taille, légèrement plus petite (planche 4*f*). Ayant des habitudes nocturnes, comme la chrysope, elle pénètre souvent dans les bâtiments éclairés, la nuit, en été. La larve ressemble à celle de la chrysope par la forme et la couleur, mais son appareil buccal est plus court et plus trapu. Les œufs allongés ne sont pas fixés à un pédoncule. Blancs au moment de la ponte, ils virent à l'orange ou au rose puis au brun foncé un peu avant l'éclosion. Au lieu d'un cocon sphérique très dense comme celui de la chrysope, l'hémérobie tisse un cocon lâche, de forme elliptique, dans un endroit abrité plutôt qu'à la vue, sur les feuilles.

L'hémérobie adulte et sa larve attaquent les petits insectes à corps mou comme les pucerons, de même que tous les stades du tétranyque. La femelle pond plus d'œufs que la chrysope et les larves de l'hémérobie sont encore plus nombreuses que celles de la chrysope dans certains endroits, au début de la saison de croissance. Le cycle biologique dure de 24 à 30 jours. On compte 23 espèces d'hémérobie au Canada.

Acariens utiles

Acariens prédateurs (Acariens : phytoséidés)

Les phytoséidés mesurent rarement plus de 0,5 mm de longueur et sont difficiles à apercevoir à l'œil nu. Avec une loupe cependant, on peut les observer sur les feuilles de fraisier et de framboisier infestées de tétranyques. Souvent globuleux ou piriformes (planche 5*a*), ils se déplacent plus rapidement que leurs congénères qui parasitent les végétaux. La différence relative de vitesse est une bonne manière de les reconnaître, mais il en existe de nombreuses espèces et l'identification exige l'œil d'un expert de même qu'un microscope sophistiqué. Une espèce, *Amblyseius fallacis* (Garman) (planche 5*a*, p. 24), se retrouve sur tous les fraisiers et framboisiers du Canada, sauf dans les provinces des Prairies.

Les acariens prédateurs se servent de leurs pièces buccales pour trouver, attraper, percer et sucer leurs proies, lesquelles comprennent des acariens, des insectes et des nématodes parasites. Beaucoup d'entre eux se nourrissent également de sève, de miellat (des pucerons), de champignons et de pollen, ce qui permet à leur population de rester stable même quand ils n'ont pas de proies. La femelle de *A. fallacis* peut pondre de 1 à 3 œufs par jour tandis que d'autres espèces en produisent quotidiennement moins d'un à plus de quatre en moyenne, selon la température et l'abondance de nourriture. Il faut de 5 à 16 jours pour que le cycle de l'œuf à l'adulte s'accomplisse. Aux températures estivales (27 °C), le cycle biologique de nombreuses espèces prédatrices dure environ 6 jours. Les phytoséidés font souvent preuve d'une grande efficacité comme agent de lutte biologique, car ils se reproduisent plus rapidement que les espèces parasites. Il est possible de s'en procurer dans le commerce au Canada où on les utilise principalement pour lutter contre le tétranyque à deux points (*Tetranychus urticae* Koch).

Autres prédateurs

Dans des conditions normales, n'importe quelle culture abritera d'autres prédateurs, souvent en grand nombre, mais il ne s'agit pas de prédateurs spécifiques aux insectes qui parasitent la culture. En font partie notamment les araignées tisseuses (planche 5*b*) comme l'épeire des jardins et les araignées fouisseuses comme les thomisides. Beaucoup de guêpes entrent également dans ce groupe, comme la guêpe jaune (planche 5c) et l'espèce *Vespula maculata* (planche 5*d*). Le rôle des asiles, qui attaquent et capturent les insectes en plein vol, n'est pas non plus à négliger. Leurs

larves, également prédatrices, détruisent les larves des coléoptères (vers blancs) ou leurs pupes sur le sol. Les libellules (planche 5*e*) et les demoiselles se font parfois nombreuses et chassent activement d'autres insectes.

Enfin, les oiseaux insectivores participent à la lutte contre les ravageurs, non seulement les insectes volants mais les chenilles et les pucerons. Quelques petits mammifères comme la taupe sont des chasseurs opportunistes qui dévorent les gros insectes ou les pupes qu'ils découvrent sur le sol ou sous terre.

Ravageurs courants du framboisier et du fraisier

Punaises et lygides (Hémiptères : miridés)

Ce groupe comprend les lygides (punaise terne) (planche 5*f*), les pentatomes et d'autres hémiptères. La plupart traversent l'hiver à l'état d'œuf, mais certains adultes hivernent sous les débris végétaux, en plein champ. Au printemps, l'œuf éclôt ou l'adulte se réveille et se met à pondre. La nymphe des lygides ressemble à un gros puceron (planche 5*f*); elle est délicate, habituellement verte et n'a pas d'ailes bien développées. Elle s'enfuit rapidement lorsqu'on la dérange. La nymphe passera par cinq stades ou mues en l'espace d'environ 5 semaines et se nourrira des sucs de la plante à chaque étape, de même qu'à l'état adulte. Le fait qu'elle se nourrit des inflorescences entraîne une pollinisation inégale à l'origine de la «face de chat» du fraisier et de la «grenaille» du framboisier. L'adulte porte des marques caractéristiques sur le dos et ses longues pièces buccales, suceuses, pointues, sont repliées sous l'abdomen. Quand on le dérange, l'insecte s'envole ou s'enfuit pour se mettre à l'abri.

Les insectes de ce groupe ne sont pas tous nuisibles, certains sont des prédateurs. Les cinq espèces qui vivent sur le fraisier s'attaquent toutes à la plante, mais l'une d'entre elles se nourrit également d'autres insectes. Parmi les 22 espèces qu'on retrouve sur le framboisier au Canada, 6 se nourrissent de la plante, 10 sont des prédateurs et 6 alternent les deux régimes.

Les lygides sont parasités par plusieurs guêpes de la famille des braconidés, au moins un ichneumon et plusieurs tachinaires. On a recensé beaucoup de prédateurs des lygides, notamment des punaises demoiselles, des punaises tête-de-clou, des miridés et des chrysopes, mais on ignore l'importance de leur rôle. Ces prédateurs préfèrent habituellement les insectes à corps mou. Toutefois, ils s'attaqueront également aux lygides, et plus particulièrement à leurs jeunes nymphes.

Pucerons (Homoptères : aphididés)

Les pucerons sont de petits insectes à corps mou qui sucent la sève des plantes. Ils sont habituellement de couleur verte, noire, jaunâtre ou rose. Durant la majeure partie de la période végétative, ils déposent leurs jeunes vivants sur la plante dont ils se nourrissent. Ce comportement crée des amas ou colonies de pucerons de tout âge (planche 6*a*, p. 28). Quand il se



Planche 6

a **Pucerons sur une feuille de fraisier (photo W. MacDiarmid)**
b **Cicadelle du genre *Macropsis* (sur un framboisier) (photo H. Severson)**
c **Charançon sur une feuille de fraisier (photo W. MacDiarmid)**
d **Anthonome de la fleur du fraisier adulte et sa larve (photo N. Bostanian)**
e **Charançon noir de la vigne adulte, sa larve et sa pupe (photo W.T. Cram)**
f **Arpenteuse du framboisier (*Autographita ampla*) (Walker)) (photo W. MacDiarmid)**



Planche 7

a **Arpenteuse du framboisier parasitée (remarquer les taches noires) (photo W. MacDiarmid)**

b **Dépouille d'arpenteuse parasitée (photo W. MacDiarmid)**

c **Rhizophage du framboisier adulte (photo W.G. Wellington)**

d **Tenthrède adulte (photo W.T. Cram)**

e **Tétranyque à deux points adulte (photo C. Chan)**

f **Femelle rouge du tétranyque à deux points, hivernant (photo C. Chan)**

nourrit, le puceron aspire tant de sève qu'il doit l'excréter rapidement, sinon il risquerait d'éclater. On appelle miellat le liquide ainsi excrété, car il renferme une grande quantité des sucs de la plante. Cette substance luisante et collante tache les végétaux près des colonies. En trop grande quantité, le miellat favorisera l'apparition d'une moisissure fuligineuse qui contaminera encore plus la plante. Le miellat attire aussi les prédateurs et les parasites des pucerons pour deux raisons : en premier lieu, beaucoup de prédateurs découvrent les pucerons en cherchant le miellat et deuxièmement, les sucres et les acides aminés que contient celui-ci constituent une bonne source d'éléments nutritifs (p. ex. pour les chrysopes adultes, les guêpes parasites, les syrphes adultes, les cécidomyies et les coccinelles).

Le puceron du framboisier, avec ses 3 mm de longueur, est passablement gros. Son corps va du vert foncé au jaune et il affectionne les jeunes pousses terminales. Le puceron du fraisier (planche 6*a*) est plus petit et vert pâle, et porte de nombreux poils. Il hiverne à l'état adulte ou sous forme de petits œufs noirs, sur les vieilles feuilles du plant. Ces deux espèces revêtent une grande importance, car elles transmettent des viroses à la plante.

Le puceron est la nourriture préférée de plusieurs insectes prédateurs comme les syrphes, les cécidomyies, les chrysopes, les coccinelles et quelques staphylins. Beaucoup d'autres insectes prédateurs, les punaises demoiselles, les réduves et les carabes par exemple, les détruisent également. Enfin, ils sont attaqués par maintes espèces de guêpes parasites, surtout celles de la famille des braconidés.

Cicadelles et cercopes (Homoptères)

Les cicadelles, qui s'attaquent au framboisier, et les cercopes, qui peuvent causer des ravages au fraisier, sont apparentés aux pucerons et détériorent les plantes un peu de la même manière, en aspirant la sève. La cicadelle est un insecte de 3 mm de longueur, très actif, au corps élancé qui fait un saut lorsqu'on le dérange. La couleur du corps varie du blanc pâle au vert brunâtre (planche 6*b*) et l'insecte se nourrit sur la face inférieure des feuilles de framboisier. Leur activité fait apparaître des taches jaunes le long des nervures et à l'extrémité des feuilles, ce qui leur donne un aspect marbré. Certaines espèces se nourrissent également des baies. La cicadelle apparaît vers la fin mai dans certaines régions, mais ailleurs on ne la voit qu'en juin et en juillet. Comme le puceron, la cicadelle excrète du miellat qui macule la plante ou le fruit d'une substance luisante et collante sur laquelle peut apparaître une moisissure fuligineuse. À l'instar des pucerons, la cicadelle est la cible de divers prédateurs comme les chrysopes, les coccinelles, certains staphylins, les punaises demoiselles, les réduves et les carabes.

On trouve des cercopes un peu partout au Canada. Quand ils se nourrissent de la sève des fraisiers, ils entraînent le nanisme de la plante et en réduisent le rendement. Cet insecte est comme recouvert de crachat, mélange d'air et d'excréments essentiel à la survie de la nymphe. L'espèce

en forme de coin qui parasite le fraisier (*Philaenus leucophthalmus* Linnaeus) a une nymphe de couleur pâle, blanchâtre à jaune, mais devient brun grisâtre à l'état adulte. L'adulte mesure 6 mm de longueur et vagabonde d'une plante à l'autre en parcourant de brèves distances par vol ou par saut. La femelle pond ses œufs orange sur la plante où la nymphe éclôt au printemps. Il n'existe pas de bon agent de lutte biologique contre les cercopes.

Charançons et autres coléoptères (Coléoptères)

Plusieurs charançons s'attaquent aux racines, à la couronne, aux bourgeons et aux feuilles du framboisier et du fraisier (planche 6*c*). Les dommages les plus importants surviennent quand la larve détruit les racines (habituellement de 5 à 15 cm sous la surface) ou la couronne. La majorité des charançons qui parasitent le fraisier et le framboisier se reproduisent de façon asexuée (sans mâle). Aucune espèce ne vole. L'adulte s'active la nuit, si bien qu'on l'aperçoit rarement durant le jour. On peut néanmoins déceler sa présence par les coches caractéristiques qui découpent la bordure des feuilles ou, dans le cas de l'anthonome de la fleur du fraisier (planche 6*d*), par les bourgeons floraux sectionnés. La larve (planches 6*d* et *e*) vit dans le sol et se nourrit des racines durant toute la période végétative. Les dommages que la larve cause à la couronne entraînent le flétrissement de la plante à la nouaison. Le plus gros charançon est le charançon noir de la vigne (*Othiorhynchus sulcatus* (Eichoff)) qui mesure 13 mm (planche 6*e*). Le plus petit, le charançon des bois (*Nemocestes incomptus* (Horn)), n'a que la moitié de cette taille.

Il est capital de différencier les charançons adultes des carabes utiles, car certains se ressemblent par la taille et la couleur. Le charançon se distingue par son «nez». En effet, la tête de l'adulte, plus étroite, termine un thorax allongé qui imite un cou; ses antennes sont couramment pliées au centre. Le corps est dur et de couleur terne et l'insecte se déplace assez lentement. Au contraire, les carabes prédateurs fuient à toute vitesse lorsqu'on les dérange. Ils ont plutôt une forme arrondie, de même qu'une carapace luisante. Ces deux types d'insectes sont nocturnes et passent la journée sous des débris ou dans des galeries.

Les charançons ne sont pas la proie d'autant d'insectes que les pucerons, mais divers vertébrés (p. ex. les taupes) de même que de plus gros carabes et des staphylins en font leurs proies. Plusieurs petits carabes mangent également les œufs des charançons. D'autres prédateurs comme les perce-oreilles s'attaquent à divers stades du charançon en laboratoire, mais on ignore leur rôle véritable dans la nature.

Divers coléoptères parasitent le framboisier et le fraisier. Le byture de la framboise existe un peu partout au Canada. L'adulte mange les jeunes feuilles et les inflorescences, ce qui réduit la nouaison. De son côté, la larve pénètre dans le fruit et provoque sa chute ou le rend impropre à la commercialisation. Ce ravageur subit les attaques des mêmes prédateurs que le charançon. La chrysomèle du fraisier est un autre coléoptère parasite qu'on trouve aussi fréquemment sur le fraisier et le framboisier.

L'adulte se nourrit des feuilles de la plante la nuit tandis que la larve en dévore les racines et la couronne. Les deux sont la cible des prédateurs qui vivent dans le sol comme les staphylins et les carabes.

Enrouleuses, chenilles, noctuelles et perceurs (Lépidoptères)

Ce groupe se compose de papillons nocturnes (planche 6*f*) dont la forme immature, une chenille, dévore différents organes de la plante : feuilles, couronne, racines et tige. Plusieurs espèces parasitent le fraisier et le framboisier à divers degrés, dans toutes les régions du Canada. Il s'agit d'un groupe diversifié de ravageurs auxquels s'attaquent des prédateurs et des parasites très variés également (tachinaires, ichneumons, braconides, chalcidies, carabes, réduves, guêpes comme la guêpe jaune et, dans une moindre mesure, les staphylins, les coccinelles, les punaises demoiselles, les anthocoris, les chrysopes et d'autres prédateurs non spécifiques comme les oiseaux et les rongeurs). En règle générale, une chenille parasitée par une guêpe ou un tachinaire porte les stigmates causés par l'inoculation des œufs (planche 7*a*, p. 29). À sa mort, elle laisse souvent une dépouille typique qu'illustrent les séquelles du parasitisme (planche 7*b*).

Quoique le rhizophage du framboisier (*Bembicia marginata* (Harris)) soit un papillon nocturne, il ressemble et se comporte comme une guêpe jaune (planche 7*c*). Ce mimétisme le met à l'abri des prédateurs comme les oiseaux qui le prennent pour une guêpe et ne s'y frôlent pas de peur de se faire piquer. Le papillon nocturne adulte volette dans les framboisiers et dépose des œufs sur la face inférieure des feuilles durant la journée, de la fin juillet à la mi-septembre. De chaque œuf émerge une petite larve jaune qui longe la ronce jusqu'au sol. Elle creuse ensuite une petite ouverture dans l'écorce et y passe l'hiver. Le printemps suivant, la larve s'enroule autour d'une nouvelle tige avant de percer une galerie jusqu'au cœur tendre et creux de la ronce qui lui servira de grenier et d'abri pour le deuxième hiver. Le printemps venu, la larve, qui a grossi, se creuse un chemin à travers la tige jusqu'à quelques centimètres au-dessus du sol. La pupaison commence immédiatement sous l'écorce. Le papillon adulte fait son apparition à la fin de juillet.

En raison de sa ressemblance avec les syrphes adultes et la guêpe jaune, le rhizophage du framboisier est rarement considéré comme un ravageur du framboisier. De même, on peut confondre les guêpes et les syrphes avec les rhizophages. Le corps du papillon est cependant recouvert d'écailles plumeuses qui lui donne une apparence veloutée et mate tandis que le corps des syrphes et de la guêpe jaune a une texture lisse et luisante.

Tenthrèdes (Hyménoptères : tenthrèdinidés)

Le tenthrède du framboisier (*Monophadnoides geniculatus* (Hartig)) adulte est une guêpe noire, aux ailes translucides, marquée de rouge et de jaune (planche 7*d*). Son corps trapu d'environ 6 mm de longueur n'a pas la taille étroite qui est typique des guêpes. Elle se manifeste en mai ou en juin et dépose ses œufs sur la feuille terminale des ronces de l'année. De l'œuf

sort une larve verte et épineuse, difficile à percevoir, car elle se confond avec le feuillage. Dans les infestations graves, la larve dévore le tissu foliaire pour ne laisser que les plus grosses nervures. À maturité, elle mesure de 12 à 17 mm de longueur et se laisse choir sur le sol où elle tisse un cocon dans lequel elle passera l'hiver. Les braconides attaquent les œufs du tenthrède, tandis que le cocon est une proie facile pour les carabes et les staphylins. Enfin la larve constitue une cible idéale pour les ichneumons et les tachinaires, de même que la guêpe jaune et les oiseaux.

Acariens

Le tétranyque à deux points (*Tetranychus urticae* Koch) adulte est difficile à voir à l'œil nu. Il s'agit d'un insecte jaune verdâtre portant deux taches foncées sur le dos (planche 7*e*). Il se nourrit en suçant la sève des plantes sur la face inférieure des feuilles. Les feuilles parasitées sont mouchetées de blanc ce qui leur donne un aspect terne et pâle. Lorsque l'infestation est grave, la feuille brunit et finit par tomber. Le tétranyque tisse une toile soyeuse sur la face inférieure des feuilles dont il se nourrit. La femelle rouge orange passe l'hiver à l'état adulte (planche 7*f*) sur les feuilles ou sous les détritus, les feuilles mortes ou le sol au pied du plant. Au printemps, elle trouve une jeune feuille dont elle se nourrira et sur laquelle elle déposera ses œufs. La population de tétranyques à deux points augmente par temps chaud et sec. Il est possible aussi que la poussière qui macule les framboisiers et les fraisiers en bordure des chemins fasse augmenter la population des tétranyques à deux points.

L'acarien du cyclamen ou du framboisier (*Phytonemus pallidus* (Banks)) est un acarien minuscule qui se nourrit dans la couronne et les feuilles du fraisier avant qu'elles se déroulent. S'il est difficile à distinguer même avec une loupe, il n'en va pas de même pour les dommages qu'il cause. En effet, les feuilles jaunissent et ne se déroulent pas; celles qui parviennent à s'ouvrir ont habituellement l'aspect du cuir et sont difformes. Les plants fortement infestés sont souvent rabougris et lorsque le ravageur s'attaque à l'inflorescence, les fruits deviennent difformes. Les acariens prédateurs comme les espèces du genre *Amblyseius* jouent un grand rôle dans la lutte contre ce parasite.

Sur la côte ouest, un ériophyide appelé phytopte du toyon (*Acalitus essigi* (Hassan)) cause d'importants ravages aux framboisiers. Cet acarien est microscopique; il passe l'hiver dans les bourgeons et dévore le fruit durant son développement au printemps et en été. Les baies touchées restent dures et sûres, même après maturité. Les acariens prédateurs permettent de combattre ce ravageur dans une certaine mesure.

Les acariens comptent beaucoup de prédateurs : chironomes, petites coccinelles noires du genre *Stethorus*, quelques petits staphylins, punaises tête-de-clou, anthocoris, chrysopes et hémérobies ainsi que d'autres acariens. Dans le sud de l'Ontario, le thrips à six taches, *Scolothrips sexmaculatus* (Pergande), s'attaque aux tétranyques. On le trouve couramment aux États-Unis, mais il n'a pas souvent été signalé au Canada.

Bibliographie

Borrer, D.J.; DeLon, D.M.; Triplehorn, C.A. 1976. An introduction to the study of insects. 4e édition. Holt Reinhart & Winston.

Carr, A. 1979. Rodale's color handbook of garden insects. Rodale Press, Erasmus, Pa.

Chant, D.A. 1985. External anatomy. Page 5–9 *in* Helle, W.; Sabelis, M.W., eds. Spider mites and their biology, natural enemies and control. Vol. 1B. Elsevier, Amsterdam.

Chant, D.A.; Hansell, R.I.C. 1971. The genus *Amblyseius* (Acari: Phytoseiidae) in Canada and Alaska. Can. J. Zool. 49:703–758.

Clausen, C.P. 1940. Entomophagous insects. McGraw Hill Biik Co., N.Y.

Clancy, D.W.; Pierce, H.D. 1966. Natural enemies of some Lygus bugs. J. Econ. Entomol. 59:853–858.

Danks, H.V. 1979. Canada and its insect fauna. Entomol. Soc. Can. Memoir 108.

Davidson, R.H.; Lyon, W.F. 1979. Insect pests of farm, garden, and orchard. 7e édition. John Wiley and Sons, N.Y.

Garth, G.S.; Shanks, C.H., Jr. 1978. Some factors affecting infestation of strawberry fields by the black vine weevil in western Washington. J. Econ. Entomol. 71:443–448.

Gordon, R.D. 1985. The Coccinellidae (Coleoptera) of America north of Mexico. J. N.Y. Entomol. Soc. 93:1–912.

Kelton, L.A. 1982. Plant bugs on fruit crops in Canada. Research Branch, Agric. Can. Monograph No. 24.

Lattin, J.D. 1989. Bionomics of the Nabidae. Ann. Rev. Entomol. 34:383–400.

Lim, K.P.; Stewart, R.K. 1976. Parasitism of the tarnished plant bug, *Lygus lineolaris* (Hempitera: Miridae) by *Peristenus pallipes* and *P. pseudopallipes* (Hymenoptera: Braconidae). Can. Entomol. 108:601–608.

McMurtry, J.A.; Hummaker, C.B.; van de Vrie, M. 1970. Ecology of tetranychid mites and their natural enemies: A review. I. Tetranychid enemies: Their biological characters and the impact of spray practices. Higardia 40:331–390.

Oatman, E.R.; Badgley, M.E.; Platner, G.R. 1985. Predators of the two-spotted spider mite on strawberry. Calif. Agric. janvier/février:9–12.

Putman, W.L. 1955. The bionomics of *Stethorus punctillum* Weise (Coleoptera: Coccinellidae) in Ontario. Can. Entomol. 87:9–33.

Schneider, F. 1969. Bionomics and physiology of aphidophagous syrphidae. Ann. Rev. Entomol. 14:103–124.

Shanks, C.H., Jr. 1981. Insects and mites. Pages 293–308 *in* Childers, N.F., ed. The strawberry. Horticultural Publications, Gainsville, Fla.

Tanigoshi, L.K. 1982. Advances in knowledge of the biology of the Phytoseiidae. Pages 1–22 *in* Hoy, M.A., ed. Recent advances in knowledge of the Phytoseiidae. Univ. Calif. Press Publ. 3284.

Ware, G.W. 1980. Complete guide to pest control—with and without chemicals. Thomson Publications, Fresno, Calif.

Publications des provinces et des États

Colombie-Britannique. 1988. Berry production guide for commercial growers. Ministère de l'Agriculture et des Pêcheries de la Colombie-Britannique, British Columbia Blueberry Co-op Association, British Columbia Cranberry Marketing Board, British Columbia Raspberry Growers Association et Fraser Valley Strawberry Growers Association.

Hunter, C.L. 1981. Insects and diseases of cane fruits. Ministère de l'Agriculture et de l'Alimentation de l'Ontario. Feuillet No. 81-001, Agdex 230.605.

Johansen, C.; Eves, J.; Retan, A. 1972. Beneficial predators and parasites found on Washington crops. Washington State University Extension, Extension Bulletin 640.

Raspberry pest identification. 1988. Plant Industry Branch, Ministère de l'Agriculture du Nouveau-Brunswick. Publ. No. 020-88, Agdex No. 232.600.

Retan, A.H.; Curtis, J. 1972*a*. Insect answers - minute pirate bug. Co-operative Extension Service, College of Agriculture, Washington State University, Pullman, E.M. 3703.

Retan, A.H.; Curtis, J. 1972*c*. Insect answers - hover or syrphid flies. Co-operative Extension Service, College of Agriculture, Washington State University, Pullman. E.M. 3704.

Strawberry pest identification. 1988. Plant Industry Branch, Ministère de l'Agriculture du Nouveau-Brunswick. Publ. No. 010-88, Agdex No. 232.600.